

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
3 June 2004 (03.06.2004)

PCT


(10) International Publication Number
WO 2004/046159 A1

- (51) International Patent Classification⁷: C07H 19/167, A61K 31/70, A61P 31/14 (74) Agent: KJELLSAA-BERGER, Hanny; Grenzacherstrasse 124, CH-4070 Basle (CH).
- (21) International Application Number: PCT/EP2003/012733
- (22) International Filing Date: 14 November 2003 (14.11.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/427,447 19 November 2002 (19.11.2002) US
60/483,970 1 July 2003 (01.07.2003) US
- (71) Applicant: F. OFFMANN-LA ROCHE AG [CH/CH]; Grenzacherstrasse 124, CH-4070 Basel (CH).
- (72) Inventors: MARTIN, Joseph, Armstrong; 350 Sharon Park Drive, Apt. I-26, Menlo Park, CA 94025 (US). SARMA, Keshab; 517 Davenport Court, Sunnyvale, CA 94087 (US). SMITH, David, Bernard; 218 West 40th Avenue, San Mateo, CA 94403 (US). SMITH, Mark; 88 Howard Street No. 808, San Francisco, CA 94105 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 2004/046159 A1

(54) Title: ANTIVIRAL NUCLEOSIDE DERIVATIVES



(57) Abstract: The present invention relates to nucleoside derivatives for the treatment of Hepatitis C viral infections including compounds of formula I (I), to pharmaceutical compositions comprising these compounds and methods for treatment or prophylaxis of Hepatitis C Virus mediated diseases employing said compounds in monotherapy or in combination therapy. The present invention further provides a process for preparing 1',3',4'-triacyl pyrimidine nucleoside from a N,1',3',4'-tetraacylpyrimidine nucleoside.

ANTIVIRAL NUCLEOSIDE DERIVATIVES

The invention relates to the field of antiviral therapy and in particular to nucleoside derivatives for treating Hepatitis C-Virus (HCV) mediated diseases. The invention provides novel chemical compounds, pharmaceutical compositions comprising these compounds, methods for treatment or prophylaxis of HCV mediated diseases employing
5 said compounds in monotherapy or in combination therapy.

The invention relates to nucleoside derivatives as inhibitors of HCV replicon RNA replication. In particular, the invention is concerned with the use of pyrimidine nucleoside compounds as inhibitors of subgenomic HCV RNA replication and pharmaceutical compositions containing such compounds.

10 Hepatitis C virus is the leading cause of chronic liver disease throughout the world. Patients infected with HCV are at risk of developing cirrhosis of the liver and subsequent hepatocellular carcinoma and hence HCV is the major indication for liver transplantation. Only two approved therapies are currently available for the treatment of HCV infection (R. G. Gish, *Sem. Liver. Dis.*, 1999 19:5). These are interferon- α
15 monotherapy and, more recently, combination therapy of the nucleoside analogue, ribavirin (Virazole), with interferon- α .


Many of the drugs approved for the treatment of viral infections are nucleosides or nucleoside analogues and most of these nucleoside analogue drugs are converted into the corresponding triphosphate *in vivo*. The triphosphates inhibit viral polymerase enzymes
20 which halts viral replication. This conversion to the triphosphate is commonly mediated by cellular kinases and therefore the direct evaluation of nucleosides as inhibitors of HCV replication is only conveniently carried out using a cell-based assay. For HCV the availability of a true cell-based viral replication assay or animal model of infection is lacking.

25 Hepatitis C virus belongs to the family of *Flaviridae*. It is an RNA virus, the RNA genome encoding a large polyprotein that after processing produces the necessary replication machinery to ensure synthesis of progeny RNA. It is believed that most of the non-structural proteins encoded by the HCV RNA genome are involved in RNA replication.

Lohmann et al. [V. Lohmann *et al.*, *Science*, 1999, 285:110-113] have described the construction of a Human Hepatoma (Huh7) cell line in which subgenomic HCV RNA molecules have been introduced and shown to replicate with high efficiency. It is believed that the mechanism of RNA replication in these cell lines is identical to the replication of the full length HCV RNA genome in infected hepatocytes. The subgenomic HCV cDNA clones used for the isolation of these cell lines have formed the basis for the development of a cell-based assay for identifying nucleoside analogue inhibitors of HCV replication.

U.S. Application Ser. No. 10/167,106 filed June 11, 2002 entitled "4'-Substituted Nucleoside Derivatives as Inhibitors of HCV RNA Replication", discloses compounds related to the present invention.

Nucleoside derivatives frequently exhibit high levels of biological activity; however, their practical utility is often limited by suboptimal physical properties and poor pharmacokinetics. The present invention relates to chemical derivatives of 4'-substituted nucleosides with improved physiochemical and pharmacokinetic properties. These derivatives more efficiently permeate the intestinal mucosa whereupon a variety of enzymes present in the cytoplasm, blood, or serum convert the derivative to the non-derivatized nucleoside. These "pronucleotides" can improve the properties such as activity, bioavailability or stability of the parent nucleotide. Administration of compounds of formula I to mammals infected by HCV inhibits subgenomic HCV replication in a hepatoma cell line.



wherein:

R^1 and R^2 are independently selected from the group consisting of hydrogen, COR^5 , $C(=O)OR^5$, $C(=O)SR^5$, $C(=O)NHR^5$ and $COCH(R^6)NHR^7$;

R^3 and R^4 independently of the other are selected from the group consisting of hydrogen, COR^5 , CO_2R^5 and $COCH(R^6)NHR^7$, or R^3 and R^4 taken together are selected from the group consisting of CH_2 , $C(CH_3)_2$ and $CHPh$;

R^5 is independently selected from the group consisting of C_{1-18} unbranched or branched alkyl, C_{1-18} unbranched or branched alkenyl, C_{1-18} unbranched or branched alkynyl, C_{1-18} lower haloalkyl, C_{3-8} cycloalkyl, alkyl substituted C_{3-8} cycloalkyl, phenyl optionally independently substituted with one to three substituents selected from the group consisting of halo, lower alkyl, lower alkoxy, lower thioalkyl, lower alkyl sulfinyl, lower alkyl sulfonyl, nitro, and cyano, CH_2Ph wherein in phenyl ring is optionally substituted as described above and CH_2OPh wherein in phenyl ring is optionally substituted as described above;

R^6 is selected from the group consisting of the side chains of naturally occurring amino acids and C_{1-5} unbranched or branched alkyl;

R^7 is selected from the group consisting of hydrogen, R^5OCO , and; hydrates, solvates, clathrates and acid addition salts thereof;

with the proviso that at least one of R^1 , R^2 , R^3 , or R^4 is other than hydrogen.

The invention provides compounds of formula I as useful agents alone or in combination with an immune system modulator, an antiviral agent, or an anti-inflammatory agent for the treatment of treating diseases mediated by Hepatitis C virus.

The invention also provides methods of treating diseases mediated by Hepatitis C virus by administering a compound of formula I. The compound can be administered alone or in combination with an immune system modulator, an antiviral agent, or an anti-inflammatory agent. The invention further includes compositions for the treatment of treating diseases mediated by Hepatitis C virus by administering a therapeutically effective amount of a compound of formula I.

Compounds of the present invention are pro-drugs or bioprecursors of the parent nucleoside and are converted *in vivo* to the compound of formula I wherein R^1 , R^2 , R^3 , and R^4 are hydrogen. Pro-drugs include acyl derivatives, amino acid esters, alkoxycarbonyl, aryloxy carbonyl, thioalkyl carbonyl and arylthiocarbonyl nucleoside or pharmaceutically acceptable salts thereof.

One embodiment of the present invention is a nucleoside derivative according to formula I wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are independently selected from the groups defined hereinabove with the proviso that at least one of R^1 , R^2 , R^3 , or R^4 is other than hydrogen.

In another embodiment of the present invention there is provided a compound according to formula I wherein R^1 , R^2 , R^3 , and R^4 each are independently COR^5 , $C(=O)OR^5$ or $C(=O)SR^5$ and each R^5 is independently selected from the group consisting of C_{1-18} unbranched or branched lower alkyl, phenyl and CH_2OPh .

- 5 In another embodiment of the present invention there is provided a compound according to formula I wherein R^1 , R^2 , R^3 , and R^4 are COR^5 and each R^5 is independently selected from the group consisting of C_{1-18} unbranched or branched lower alkyl, phenyl and CH_2OPh .

- 10 In another embodiment of the present invention there is provided a compound according to formula I wherein R^1 is COR^5 , $C(=O)OR^5$, $C(=O)SR^5$ or $COCH(R^6)NHR^7$; R^2 , R^3 and R^4 are hydrogen; and, R^5 or R^6 and R^7 are as defined hereinabove.

- In another embodiment of the present invention there is provided a compound according to formula I wherein R^1 is COR^5 , $C(=O)OR^5$, $C(=O)SR^5$ or $COCH(R^6)NHR^7$; R^2 , R^3 and R^4 are hydrogen; R^5 is selected from a group consisting of C_{1-18} unbranched or
15 branched lower alkyl, C_{3-8} cycloalkyl, phenyl and CH_2OPh or R^6 is selected from the group consisting of C_{1-5} unbranched or branched alkyl and the side chain of a naturally occurring amino acid and R^7 is as defined hereinabove.

- In another embodiment of the present invention there is provided a compound according to formula I wherein R^2 is selected from the group consisting of COR^5 ,
20 $C(=O)OR^5$, $C(=O)SR^5$ and $COCH(R^6)NHR^7$; R^1 , R^3 and R^4 are hydrogen; and R^5 or R^6 and R^7 are as defined hereinabove.

- In another embodiment of the present invention there is provided a compound according to formula I wherein R^2 is selected from the group consisting of COR^5 , $C(=O)OR^5$, $C(=O)SR^5$ and $COCH(R^6)NHR^7$; R^1 , R^3 and R^4 are hydrogen; R^5 is selected
25 from the group consisting of C_{1-18} unbranched or branched alkyl, C_{3-8} cycloalkyl and phenyl or R^6 is C_{1-5} unbranched or branched alkyl or the side chain of a naturally occurring amino acid and R^7 are as defined hereinabove.

- In another embodiment of the present invention there is provided a compound according to formula I wherein R^2 is $COCH(R^6)NH_2$; R^1 , R^3 , R^4 and R^7 are hydrogen;
30 and, R^6 is selected from the group consisting of C_{1-5} unbranched or branched alkyl or CH_2Ph .

In another embodiment of the present invention there is provided a compound according to formula I wherein R^1 and R^2 are independently selected from the group consisting of COR^5 , $C(=O)OR^5$, $C(=O)SR^5$ and $COCH(R^6)NHR^7$; R^3 and R^4 are hydrogen; and, R^5 and/or R^6 and R^7 are independently selected from the group defined
5 hereinabove.

In another embodiment of the present invention there is provided a compound according to formula I wherein R^1 is hydrogen; R^2 , R^3 and R^4 are selected from the group consisting of COR^5 , $C(=O)OR^5$ or $C(=O)SR^5$; and R^5 is independently selected from the group defined hereinabove.

10 In another embodiment of the present invention there is provided a compound wherein R^1 is hydrogen; R^2 is COR^5 , $C(=O)OR^5$, $C(=O)SR^5$ or $COCH(R^6)NHR^7$; R^3 and R^4 taken together are selected from the group consisting of CH_2 , $C(CH_3)_2$ and $CHPh$; and, R^5 or R^6 and R^7 are independently selected from the group hereinabove.

In another embodiment of the present invention there is provided a compound
15 according to formula I wherein R^1 and R^2 are hydrogen; R^3 and R^4 taken independently are selected from the group consisting of COR^5 , $C(=O)OR^5$, $C(=O)SR^5$ and $COCH(R^6)NHR^7$; and, R^5 or R^6 and R^7 are independently selected from the group hereinabove.

In another embodiment of the present invention there is provided a compound
20 according to formula I wherein R^1 and R^2 are selected from the group consisting of COR^5 , $C(=O)OR^5$, $C(=O)SR^5$ and $COCH(R^6)NHR^7$; R^3 and R^4 taken together are selected from the group consisting of CH_2 , $C(CH_3)_2$ and $CHPh$; R^7 is hydrogen or $C(=O)OR^5$; and R^5 and/or R^6 are independently selected from the group defined hereinabove.

The another embodiment of the present invention there is provided a compound
25 according to formula I wherein R^1 and R^2 are selected from the group consisting of hydrogen, COR^5 , $C(=O)OR^5$ and $COCH(R^6)NHR^7$; R^3 and R^4 independently of the other are selected from the group consisting of hydrogen, COR^5 , CO_2R^5 and $COCH(R^6)NHR^7$, or R^3 and R^4 taken together are selected from the group consisting of CH_2 , $C(CH_3)_2$ and $CHPh$; R^5 is independently selected from the group consisting of C_{1-6} unbranched or
30 branched alkyl, C_{1-6} unbranched or branched alkenyl, C_{1-6} unbranched or branched alkynyl, C_{1-6} lower haloalkyl, C_{3-8} cycloalkyl, alkyl substituted C_{3-8} cycloalkyl, phenyl optionally independently substituted with one to three substituents selected from the group consisting of halo, lower alkyl, lower alkoxy, lower thioalkyl, lower alkyl sulfinyl,

lower alkyl sulfonyl, nitro, and cyano, CH_2Ph wherein in phenyl ring is optionally substituted as described above and CH_2OPh wherein in phenyl ring is optionally substituted as described above; R^6 and R^7 are as defined above, with the proviso that at least one of R^1 , R^2 , R^3 , or R^4 is other than hydrogen.

- 5 According to the present invention nucleoside derivatives according to formula I wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove are useful agents in the control of diseases mediated by HCV.

- According to the present invention compounds according to formula I wherein R^1 , R^2 , R^3 , or R^4 are each independently selected from the group consisting of COR^5 , $\text{C}(=\text{O})\text{OR}^5$ and $\text{C}(=\text{O})\text{SR}^5$ and R^5 are independently selected from the group defined hereinabove are useful agents in the control of diseases mediated by HCV.
- 10

- According to the present invention compounds according to formula I wherein R^1 , R^2 , R^3 , or R^4 are each COR^5 and each R^5 is independently selected from the group consisting of C_{1-18} unbranched or branched alkyl, phenyl and CH_2OPh are useful agents in the control of diseases mediated by HCV.
- 15

According to the present invention compounds according to formula I wherein R^1 is selected from the group consisting of COR^5 , CO_2R^5 , $\text{C}(=\text{O})\text{SR}^5$ and $\text{COCH}(\text{R}^6)\text{NHR}^7$; R^2 , R^3 , and R^4 are each hydrogen; and, R^5 or R^6 and R^7 are as defined hereinabove are useful agents in the control of diseases mediated by HCV.

- 20 According to the present invention compounds according to formula I wherein R^1 is selected from the group consisting of COR^5 , $\text{C}(=\text{O})\text{OR}^5$, $\text{C}(=\text{O})\text{SR}^5$ and $\text{COCH}(\text{R}^6)\text{NHR}^7$; R^2 , R^3 , and R^4 are each hydrogen; R^5 is selected from a group consisting of C_{1-18} unbranched or branched lower alkyl, C_{3-8} cycloalkyl, phenyl and CH_2OPh , or R^6 is selected from the group consisting of C_{1-5} unbranched or branched alkyl and the side chain of a naturally occurring amino acid and R^7 is hydrogen are useful agents in the control of diseases mediated by HCV.
- 25

- According to the present invention compounds according to formula I wherein R^2 is selected from the group consisting of COR^5 , $\text{C}(=\text{O})\text{OR}^5$, $\text{C}(=\text{O})\text{SR}^5$ and $\text{COCH}(\text{R}^6)\text{NHR}^7$; R^1 , R^3 and R^4 are hydrogen; and, R^5 or R^6 and R^7 are as defined hereinabove are useful agents in the control of diseases mediated by HCV.
- 30

According to the present invention compounds according to formula I wherein R^2 is selected from the group consisting of COR^5 , CO_2R^5 , and $\text{COCH}(\text{R}^6)\text{NHR}^7$; R^1 , R^3 and R^4

are hydrogen; R^5 is selected from a group consisting of C_{1-18} unbranched or branched alkyl, C_{3-8} cycloalkyl, phenyl and CH_2OPh or R^6 is C_{1-5} unbranched or branched alkyl or the side chain of a naturally occurring amino acid and R^7 is hydrogen are useful agents in the control of diseases mediated by HCV.

- 5 According to the present invention compounds according to formula I wherein R^2 is $COCH(R^6)NHR^7$; R^1 , R^3 and R^4 are hydrogen; R^6 is selected from the group consisting of C_{1-5} unbranched or branched alkyl and CH_2Ph ; and, R^7 is hydrogen are useful agents in the control of diseases mediated by HCV.

- 10 According to the present invention compounds according to formula I wherein R^3 and R^4 are hydrogen; and, R^1 , R^2 , R^5 , R^6 and R^7 are independently selected from the groups defined hereinabove are useful agents in the control of diseases mediated by HCV.

- 15 According to the present invention compounds according to formula I wherein R^1 is hydrogen; R^2 , R^3 and R^4 are independently selected from the group consisting of COR^5 , $C(=O)OR^5$ and $C(=O)SR^5$ and, each R^5 is independently selected from the group defined hereinabove are useful agents in the control of diseases mediated by HCV.

- 20 According to the present invention compounds according to formula I wherein R^1 is hydrogen; R^3 and R^4 , taken together, are selected from the group consisting of CH_2 , $C(CH_3)_2$ and $CHPh$; R^2 , R^5 or R^6 and R^7 are as defined hereinabove are useful agents in the control of diseases mediated by HCV.

According to the present invention compounds according to formula I wherein R^1 and R^2 are hydrogen; R^3 , R^4 , R^5 and/or R^6 and R^7 are independently selected from the group defined hereinabove are useful agents in the control of diseases mediated by HCV.

- 25 According to the present invention compounds according to formula I wherein R^3 and R^4 taken together are selected from the group consisting of CH_2 , $C(CH_3)_2$ and $CHPh$; R^1 , R^2 , R^5 and/or R^6 and R^7 are independently selected from the group defined hereinabove are useful agents in the control of diseases mediated by HCV.

According to the present invention compounds according to formula I, wherein R^1 to R^7 are as defined hereinabove are useful agents in the control of diseases mediated by HCV.

- 30 According to the present invention compounds according to formula I, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in combination with at least one

immune system modulator and/or antiviral agent that inhibits replication of HCV are useful agents in the control of diseases mediated by HCV.

According to the present invention compounds according to formula I, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in combination with at least one
5 immune system modulator are useful agents in the control of diseases mediated by HCV.

According to the present invention compounds according to formula I, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in combination with an interferon, interleukin, tumor necrosis factor, colony stimulating factor or an anti-inflammatory agent are useful agents in the control of diseases mediated by HCV.

10

According to the present invention compounds according to formula I, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in combination an interferon are useful agents in the control of diseases mediated by HCV.

According to the present invention compounds according to formula I, wherein R^1 , R^2 ,
15 R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in combination with interferon- α or a chemically derivatized interferon are useful agents in the control of diseases mediated by HCV.

According to the present invention compounds according to formula I, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in combination with at least one other
20 anti-viral agent are useful agents in the control of diseases mediated by HCV.

According to the present invention compounds according to formula I, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in combination with at least one HCV protease inhibitor, HCV polymerase inhibitor, HCV helicase inhibitor, HCV primase inhibitor, HCV integrase inhibitor or HCV fusion inhibitor are useful agents in the
25 control of diseases mediated by HCV.

In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a nucleoside derivatives according to formula I wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove.

30 In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I wherein R^1 , R^2 ,

R^3 , or R^4 are each independently selected from the group consisting of COR^5 , $C(=O)OR^5$ and $C(=O)SR^5$ and R^5 are independently selected from the group defined hereinabove.

In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a
5 therapeutically effective quantity of a compound according to formula I wherein R^1 , R^2 , R^3 , or R^4 are each COR^5 and each R^5 is independently selected from the group consisting of C_{1-18} unbranched or branched alkyl, phenyl and CH_2OPh .

In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a
10 therapeutically effective quantity of a compound according to formula I wherein R^1 is selected from the group consisting of COR^5 , CO_2R^5 , $C(=O)SR^5$ and $COCH(R^6)NHR^7$; R^2 , R^3 , and R^4 are each hydrogen; and, R^5 or R^6 and R^7 are as defined hereinabove.

In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a
15 therapeutically effective quantity of a compound according to formula I wherein R^1 is selected from the group consisting of COR^5 , $C(=O)OR^5$, $C(=O)SR^5$ and $COCH(R^6)NHR^7$; R^2 , R^3 , and R^4 are each hydrogen; R^5 is selected from a group consisting of C_{1-18} unbranched or branched lower alkyl, C_{3-8} cycloalkyl, phenyl and CH_2OPh , or R^6 is selected from the group consisting of C_{1-5} unbranched or branched alkyl and the side
20 chain of a naturally occurring amino acid and R^7 is hydrogen.

In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I wherein R^2 is selected from the group consisting of COR^5 , $C(=O)OR^5$, $C(=O)SR^5$ and
25 $COCH(R^6)NHR^7$; R^1 , R^3 and R^4 are hydrogen; and, R^5 or R^6 and R^7 are as defined hereinabove.

In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I wherein R^2 is
30 selected from the group consisting of COR^5 , CO_2R^5 , $C(=O)SR^5$ and $COCH(R^6)NHR^7$; R^1 , R^3 and R^4 are hydrogen; R^5 is selected from a group consisting of C_{1-18} unbranched or branched alkyl, C_{3-8} cycloalkyl, phenyl and CH_2OPh or R^6 is C_{1-5} unbranched or branched alkyl or the side chain of a naturally occurring amino acid and R^7 is hydrogen.

In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I wherein R^2 is $\text{COCH}(R^6)\text{NHR}^7$; R^1 , R^3 and R^4 are hydrogen; R^6 is selected from the group consisting of C_{1-5} unbranched or branched alkyl and CH_2Ph ; and, R^7 is hydrogen.

In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I wherein R^3 and R^4 are hydrogen; and, R^1 , R^2 , R^5 , R^6 and R^7 are independently selected from the groups defined hereinabove.

In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I wherein R^1 is hydrogen; R^2 , R^3 and R^4 are independently selected from the group consisting of COR^5 , $\text{C}(=\text{O})\text{OR}^5$ and $\text{C}(=\text{O})\text{SR}^5$ and, each R^5 is independently selected from the group defined hereinabove.

In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I wherein R^1 is hydrogen; R^3 and R^4 , taken together, are selected from the group consisting of CH_2 , $\text{C}(\text{CH}_3)_2$ and CHPh ; R^2 , R^5 or R^6 and R^7 are as defined hereinabove.

In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I wherein R^1 and R^2 are hydrogen; R^3 , R^4 , R^5 and/or R^6 and R^7 are independently selected from the group defined hereinabove.

In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I wherein R^3 and R^4 taken together are selected from the group consisting of CH_2 , $\text{C}(\text{CH}_3)_2$ and CHPh ; R^1 , R^2 , R^5 and/or R^6 and R^7 are independently selected from the group defined hereinabove.

In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a

therapeutically effective quantity of a compound according to formula I, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in a dose between 1 and 100 mg/kg of body weight per day.

5 In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a human in need thereof a therapeutically effective quantity of a compound according to formula I, wherein R^1 to R^7 are as defined hereinabove.

10 In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in combination with at least one immune system modulator and/or antiviral agent that inhibits replication of HCV.

15 In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in combination with at least one immune system modulator.

20 In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in combination with an interferon, interleukin, tumor necrosis factor, colony stimulating factor or an anti-inflammatory agent.

25 In another embodiment of the present invention there is provided a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in combination an interferon.


30 In another embodiment of the present invention is a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in combination with interferon- α or a chemically derivatized interferon.

In another embodiment of the present invention is a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in combination with at least one other
5 anti-viral agent.

In another embodiment of the present invention is a method for treating diseases mediated by HCV comprising administering to a mammal in need thereof a therapeutically effective quantity of a compound according to formula I, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in combination with at least one HCV
10 protease inhibitor, HCV polymerase inhibitor, HCV helicase inhibitor, HCV primase inhibitor, HCV integrase inhibitor or HCV fusion inhibitor.

In another embodiment of the present invention is a pharmaceutical composition comprising a therapeutically effective quantity of a compound of formula I, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined hereinabove, in combination with one or more
15 pharmaceutically acceptable carrier, diluent or excipient with the proviso that at least one of R^1 , R^2 , R^3 , or R^4 is other than hydrogen.

In another embodiment of the present invention there is provided a process for converting an N-acyl cytidine compound IVa to a cytidine compound IVb by selective
20 cleavage of an N-acyl moiety from IVa



wherein R^2 , R^3 , R^4 and R^5 are as defined hereinabove, said process comprising contacting a solution of said N-acyl pyrimidine nucleoside with ZnBr_2 in a protic solvent R^aOH wherein R^a is hydrogen or C_{1-4} alkyl.

25 In another embodiment of the present invention there is provided a process for converting IVa to IVb wherein R^2 , R^3 , R^4 and R^5 are as defined hereinabove, said process

comprising contacting a solution of said N-acyl pyrimidine nucleoside with ZnBr_2 in methanol and optimally with an aprotic organic solvent.

The phrase "a" or "an" entity as used herein refers to one or more of that entity; for example, a compound refers to one or more compounds or at least one compound. As
5 such, the terms "a" (or "an"), "one or more", and "at least one" can be used interchangeably herein.

The phrase "as defined hereinabove" refers to the first definition for each group as provided in the definition of formula I.

The terms "optional" or "optionally" as used herein means that a described event or
10 circumstance may or may not occur, and that the description includes instances where said event or circumstance occurs and instances in which it does not. For example, "optionally substituted phenyl" means that the phenyl may or may not be substituted and that the description includes both unsubstituted phenyl and phenyl wherein there is substitution.

15 Compounds of the present invention may have asymmetric centers located on the side chain of a carboxylic ester, amide or carbonate moiety that produce diastereomers when linked to the nucleoside. All stereoisomers on the side chain of the compounds of the instant invention are contemplated, either in admixture or in pure or substantially pure
20 form. The definition of the compounds according to the invention embraces all possible stereoisomers and their mixtures. It also embraces the racemic forms as well as the isolated optical isomers. The racemic forms can be resolved by physical methods, such as, for example, fractional crystallization, separation or crystallization of diastereomeric derivatives or separation by chiral column chromatography. The individual optical
25 isomers can be obtained from the racemates by conventional methods, such as, for example, salt formation with an optically active acid followed by crystallization.

All configurational isomers of compounds of the present invention are contemplated, either in admixture or in pure or substantially pure form. The definition of compounds of the present invention embraces both *cis* and *trans* isomers of cycloalkyl rings.

30 The term "alkyl" as used herein denotes an unbranched or branched chain hydrocarbon residue containing 1 to 18 carbon atoms. Preferred alkyl groups are unbranched or branched hydrocarbon residues containing 1 to 12 carbon atoms. The term "lower alkyl" denotes an unbranched or branched chain hydrocarbon residue containing 1 to 6 carbon

atoms. Representative lower alkyl groups include methyl, ethyl, propyl, *i*-propyl, *n*-butyl, *i*-butyl, *t*-butyl or pentyl.

The term "haloalkyl" as used herein denotes an unbranched or branched chain alkyl group as defined above wherein 1, 2, 3 or more hydrogen atoms are substituted by a
5 halogen. Examples are 1-fluoromethyl, 1-chloromethyl, 1-bromomethyl, 1-iodomethyl, trifluoromethyl, trichloromethyl, tribromomethyl, triiodomethyl, 1-fluoroethyl, 1-chloroethyl, 1-bromoethyl, 1-iodoethyl, 2-fluoroethyl, 2-chloroethyl, 2-bromoethyl, 2-iodoethyl, 2,2-dichloroethyl, 3-bromopropyl or 2,2,2-trifluoroethyl.

The term "cycloalkyl" as used herein denotes a saturated carbocyclic ring containing 3 to
10 8 carbon atoms, i.e. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl.

The term "alkenyl" as used herein denotes an unsubstituted [or substituted] hydrocarbon chain radical having from 2 to 18 carbon atoms, preferably 2 to 7 and especially preferred from 2 to 4 carbon atoms, and having one or two olefinic double bonds, preferably one
15 olefinic double bond. Examples are vinyl, 1-propenyl, 2-propenyl (allyl) or 2-butenyl (crotyl).

The term "alkynyl" as used herein denotes an unsubstituted hydrocarbon chain radical having from 2 to 18 carbon atoms, preferably 2 to 7 and especially preferred from 2 to 4 carbon atoms, and having one or where possible two triple bonds[, preferably one triple
20 bond]. Examples are ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl or 3-butylnyl.

The term "alkoxy" as used herein denotes an unsubstituted unbranched or branched chain alkyloxy group wherein the "alkyl" portion is as defined above such as methoxy, ethoxy, *n*-propyloxy, *i*-propyloxy, *n*-butyloxy, *i*-butyloxy, *t*-butyloxy, pentyloxy, hexyloxy, heptyloxy including their isomers. "Lower alkoxy" as used herein denotes an
25 alkoxy group with a "lower alkyl" group as previously defined.

The term "alkylthio" or "thioalkyl" as used herein denotes a unbranched or branched chain (alkyl)S- group wherein the "alkyl" portion is as defined above. Examples are methylthio, ethylthio, *n*-propylthio, *i*-propylthio, *n*-butylthio, *i*-butylthio or *t*-butylthio.

The term "alkoxyalkyl" as used herein denotes an alkoxy group as defined above which is
30 bonded to an alkyl group as defined above. Examples are methoxymethyl, methoxyethyl, methoxypropyl, ethoxymethyl, ethoxyethyl, ethoxypropyl, propyloxypropyl,

methoxybutyl, ethoxybutyl, propyloxybutyl, butyloxybutyl, *t*-butyloxybutyl, methoxypentyl, ethoxypentyl, and propyloxypentyl including their isomers.

The term "hydroxyalkyl" as used herein denotes a unbranched or branched chain alkyl group as defined above wherein 1, 2, 3 or more hydrogen atoms are substituted by a hydroxy group. Examples are hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 1-hydroxypropyl, 2-hydroxypropyl, 3-hydroxypropyl, hydroxyisopropyl, hydroxybutyl and the like.

The term "aryl" as used herein denotes an optionally substituted monocyclic or polycyclic-aromatic group comprising carbon and hydrogen atoms. Examples of suitable aryl groups include, but are not limited to, phenyl and naphthyl (e. g. 1-naphthyl or 2-naphthyl). Suitable substituents for aryl are selected from the group consisting of alkyl, alkenyl, alkynyl, aryloxy, cycloalkyl, acyl, acylamino, alkoxy, amino, alkylamino, dialkylamino, halogen, haloalkyl, hydroxy, nitro and cyano.

The term "acyl" ("alkylcarbonyl") as used herein denotes a group of formula $C(=O)R$ wherein R is hydrogen, unbranched or branched alkyl containing 1 to 7 carbon atoms or a phenyl group.

The terms "alkoxycarbonyl" and "aryloxycarbonyl" as used herein denotes a group of formula $-C(=O)OR$ wherein R is alkyl or aryl respectively and alkyl and aryl are as defined herein.

The terms "thioalkylcarbonyl" and "arylthiocarbonyl" as used herein denotes a group of formula $-C(=O)SR$ wherein R is alkyl or aryl respectively and alkyl and aryl are as defined herein.



The term halogen stands for fluorine, chlorine, bromine or iodine, preferably fluorine, chlorine, bromine.

The term "amino acid" as used herein refers to naturally occurring amino acids, as well as to optical isomers (enantiomers and diastereomers), synthetic analogs and derivatives thereof. α -Amino acids comprise a carbon atom bonded to a carboxyl group, an amino group, a hydrogen atom and a unique "side chain" group. The term "naturally occurring amino acids" means the L-isomers of the naturally occurring amino acids. The naturally occurring amino acids are glycine, alanine, valine, leucine, isoleucine, serine, methionine, threonine, phenylalanine, tyrosine, tryptophan, cysteine, proline, histidine, aspartic acid, asparagine, glutamic acid, glutamine, γ -carboxyglutamic acid, arginine, ornithine and

lysine. The side chains of naturally occurring amino acids include: hydrogen, methyl, *iso*-propyl, *iso*-butyl, *sec*-butyl, -CH₂OH, -CH(OH)CH₃, -CH₂SH, -CH₂CH₂SMe, -
 (CH₂)_pCOR wherein R is -OH or -NH₂ and p is 1 or 2, -(CH₂)_q-NH₂ where q is 3 or 4, -
 (CH₂)₃-NHC(=NH)NH₂, -CH₂C₆H₅, -CH₂-*p*-C₆H₄-OH, (3-indolyl)methylene, (4-
 5 imidazolyl)methylene.

The term "acylating agent" as used herein refers to either an anhydride, acyl halide or other activated derivative of a carboxylic acid. The term "anhydride" as used herein refers to compounds of the general structure RC(O)-O-C(O)R wherein is as defined in the previous paragraph. The term "acyl halide" as used herein refers to the group RC(O)X
 10 wherein X is bromo or chloro. The term "activated derivative" of a compound as used herein refers to a transient reactive form of the original compound which renders the compound active in a desired chemical reaction, in which the original compound is only moderately reactive or non-reactive. Activation is achieved by formation of a derivative or a chemical grouping within the molecule with a higher free energy content than that of
 15 the original compound, which renders the activated form more susceptible to react with another reagent. In the context of the present invention activation of the carboxy group is of particular importance. The term acylating agent as used herein further includes reagents that produce carbonates (-OC(=O)OR⁵), carbamates (-NHC(=O)OR⁵), thiocarbonate(-OC(=O)SR⁵), and thiocarbamate (-NHC(=O)SR⁵), derivatives such as
 20 alkoxychlorocarbonates, R⁵OC(=O)Cl, and alkylthiochlorocarbonates, R⁵SC(=O)Cl, wherein R⁵ is as defined hereinabove.

The term "protecting group" as used herein means a chemical group that (a) preserves a reactive group from participating in an undesirable chemical reaction; and (b) can be easily removed after protection of the reactive group is no longer required. For example,
 25 the trialkylsilyl is a protecting group for a primary hydroxyl function and an acetonide is a protecting group for a *vicinal* diol.

In the pictorial representation of the compounds given throughout this application, a thickened tapered line () indicates a substituent which is above the plane of the ring to which the asymmetric carbon belongs and a dotted line () indicates a
 30 substituent which is below the plane of the ring to which the asymmetric carbon belongs.

The term "combination" as used herein in reference in administering a plurality of drugs in a therapeutic regimen by concurrent or sequential administration of the drugs at the same time or at different times.

The term "chemically-derivatized interferon" as used herein refers to an interferon molecule covalently linked to a polymer which alters the physical and/or pharmacokinetic properties of the interferon. A non-limiting list of such polymers include polyalkylene oxide homopolymers such as polyethylene glycol (PEG) or polypropylene glycol (PPG),
5 polyoxyethylenated polyols, copolymers thereof and block copolymers thereof, provided that the water solubility of the block copolymers is maintained. One skilled in the art will be aware of numerous approaches to linking the polymer and interferon (for example, see A. Kozlowski and J. M. Harris *J. Control. Release* 2001 72(1-3):217-24). A non-limiting list of chemically derivatized IFN α contemplated in the present patent include peginterferon-
10 α -2a (PEGASYS®) and peginterferon- α -2b (PEGINTRON®).

Compounds of formula I exhibit tautomerism. Tautomeric compounds can exist as two or more interconvertable species. In many cases these entities result from the shift a covalently bonded hydrogen atom shift between two atoms. Tautomeric compounds exist in equilibrium with each other, so that attempts to isolate the individual tautomers
15 usually produce a mixture having chemical and physical properties consistent with a mixture of compounds. The position of the equilibrium is dependent on chemical features within the molecule. For example, in many aliphatic aldehydes and ketones, such as acetaldehyde, the keto form predominates while; in phenols, the enol form predominates. The most common type of tautomerism is that involving carbonyl, (or
20 keto) compounds and vinyl alcohols (or enols which arise from a hydrogen atom shift between the carbon and oxygen atoms and a concomitant shift in the position of the double bond. The present invention includes lactams which can exist as amide or hydroxy substituted heterocyclic forms.

Compounds of formula I which are basic can form pharmaceutically acceptable salts with
25 inorganic acids such as hydrohalic acids (e.g. hydrochloric acid and hydrobromic acid), sulphuric acid, nitric acid and phosphoric acid, and the like, and with organic acids (e.g. with acetic acid, tartaric acid, succinic acid, fumaric acid, maleic acid, malic acid, salicylic acid, citric acid, methanesulphonic acid and *p*-toluene sulphonic acid, and the like). The formation and isolation of such salts can be carried out according to methods known in
30 the art.

The term "solvate" as used herein means a compound of the invention or a salt, thereof, that further includes a stoichiometric or non-stoichiometric amount of a solvent bound by non-covalent intermolecular forces. Preferred solvents are volatile, non-toxic, and/or acceptable for administration to humans in trace amounts.

The term "hydrate" as used herein means a compound of the invention or a salt thereof, that further includes a stoichiometric or non-stoichiometric amount of water bound by non-covalent intermolecular forces.

The term "clathrate" as used herein means a compound of the invention or a salt thereof in the form of a crystal lattice that contains spaces (e.g., channels) that have a guest molecule (e.g., a solvent or water) trapped within.

In general, the nomenclature used in this Application and given in Table 1-A is based on AUTONOMTM v.4.0, a Beilstein Institute computerized system for the generation of IUPAC systematic nomenclature.

Examples of representative compounds within the scope of the invention are provided in the following table 1 and 1-A. These examples and preparations are provided to enable those skilled in the art to more clearly understand and to practice the present invention. They should not be considered as limiting the scope of the invention, but merely as being illustrative and representative thereof. The compounds of formula I may be prepared by various methods known in the art of organic chemistry.


TABLE 1							
Cpd No.	 (I)				Method	Mass Spectrum (M+H) ⁺	Melting Point
	R ¹	R ²	R ³	R ⁴			
1	MeCO	MeCO	MeCO	MeCO	A	453	77.0-80.1
2	EtCO	EtCO	EtCO	EtCO	A	508 (M) ⁺	
3	<i>n</i> -PrCO	<i>n</i> -PrCO	<i>n</i> -PrCO	<i>n</i> -PrCO	A	564 (M) ⁺	
4	<i>i</i> -BuCO	<i>i</i> -BuCO	<i>i</i> -BuCO	<i>i</i> -BuCO	A	621	
5	<i>t</i> -BuCO	<i>t</i> -BuCO	<i>t</i> -BuCO	<i>t</i> -BuCO	A	621	67.4-80.0
6	PhCO	PhCO	PhCO	PhCO	A	700 (M) ⁺	106.9-116.1
7	PhOCH ₂ CO	PhOCH ₂ CO	PhOCH ₂ CO	PhOCH ₂ CO	A	821	-
8	<i>t</i> -BuOCO	H	H	H	B	385	80.0-83.5
9	MeCO	H	H	H	B	327	94.0-96.8
10	EtCO	H	H	H	B	341	
11	<i>n</i> -PrCO	H	H	H	B	355	
12	<i>i</i> -PrCO	H	H	H	B	355	
13	<i>t</i> -BuCO	H	H	H	B	369	108.0-114.0
14	PhCO	H	H	H	B	389	108.1-118.2
15	PhCH ₂ CO	H	H	H	B	403	
16	PhOCH ₂ CO	H	H	H	B	419	
17	<i>n</i> -BuOCO	H	H	H	B	385	86.5-94.0
18	H	Val HCl	H	H	D	384	
19	H	Phe HCl	H	H	D	432	
20	H	Ala HCl	H	H	D	356	

TABLE 1


Cpd No.	 (I)				Method	Mass Spectrum (M+H)+	Melting Point
	R ¹	R ²	R ³	R ⁴			
21	H	PhCO	H	H	D	389	
22	H	MeCO	MeCO	MeCO	C	411	
23	H	EtCO	EtCO	EtCO	C	453	75.9-79.9
24	H	<i>i</i> -PrCO	<i>i</i> -PrCO	<i>i</i> -PrCO	C	495	
25	H	PhCO	PhCO	PhCO	C		
26	H	PhCO	PhCO	3-Cl-PhCO	C		
27	<i>n</i> -BuOCO	PhCO	PhCO	PhCO	C	697	
28	PhCH ₂ CO	<i>i</i> -PrCO	H	H	E	473	
29	H	<i>n</i> -PrCO	C(CH ₃) ₃		D	395	
30	H	PhCO	C(CH ₃) ₃		D	429	
31	H	Boc-Phe	C(CH ₃) ₃		D	572	
32	H	<i>i</i> -PrCO	C(CH ₃) ₃		D	395	
33	H	H	C(CH ₃) ₃		-	325	106.2-120.1
34	<i>i</i> -PrCO	<i>i</i> -PrCO	C(CH ₃) ₃		E	465	
35	MeCO	MeCO	C(CH ₃) ₃		E	409	105.0-109.9
36	PhCO	PhCO	C(CH ₃) ₃		E	533	
37	MeOCO	MeOCO	MeOCO	MeOCO	A	517	
38 ¹	Val-NH-Boc	H	H	H	A	484	120.2-121.3
39	H	<i>i</i> -PrCO	H	H	D	355	
40 ²	H	<i>i</i> -PrCO	H	H	D	355	
41 ²	H	<i>n</i> -PrCO	<i>n</i> -PrCO	<i>n</i> -PrCO	C	495	52.6-58.4
42	C ₈ H ₁₇ OCO	H	H	H	B	441	
43 ²	H	PhCO	H	H	D	389	163-166.5
44	C ₇ H ₁₅ OCO	H	H	H	B	427	
45 ²	H	<i>i</i> -BuCO	<i>i</i> -BuCO	<i>i</i> -BuCO	C	537	142-142.8
46 ²	H	<i>n</i> -BuCO	<i>n</i> -BuCO	<i>n</i> -BuCO	C		72.9-74.7
47	C ₆ H ₁₃ OCO	H	H	H	B	413	134.4-136.0
48	H	H	<i>i</i> -PrCO	<i>i</i> -PrCO	F	425	
49 ²	H	Ile HCl	H	H	D	398	
50 ²	H	<i>t</i> -BuCO	<i>t</i> -BuCO	<i>t</i> -BuCO	F		136.2-140
51	<i>n</i> -PrOCO	H	H	H	B	371	
52	<i>n</i> -C ₅ H ₁₁ OCO	H	H	H	B	399	
53	<i>i</i> -BuOCO	H	H	H	B	385	
54 ²	H	MeCO	MeCO	MeCO	C	411	88.0-90.9
55	<i>i</i> -PrCO	<i>i</i> -PrCO	<i>i</i> -PrCO	<i>i</i> -PrCO	A	565	60.4-64
56 ²	H	<i>i</i> -PrCO	<i>i</i> -PrCO	<i>i</i> -PrCO	A		145-146.9
57	MeOCO	H	H	H	B		141.1-141.8
58	EtOCO	H	H	H	B	357	
59 ²	H	EtCO	EtCO	EtCO	C		72-75.2
60	PhCO	EtCO	EtCO	EtCO	C ⁴	557	54.6-56.9
61 ²	H	<i>n</i> -PrCO	H	H	D	355	
62 ²	H	PhCO	PhCO	PhCO	D	597	208.9-210.3
63 ²	H	<i>n</i> -C ₅ H ₁₁ CO	H	H	D	383	
64 ³	H	COOC ₆ H ₁₃	H	H	D		71.0-103.9
65 ³	H	COOC ₇ H ₁₅	H	H	D		90.9-94.6
66 ³	H	COOC ₈ H ₁₇	H	H	D		70.5-81
67 ²	H	<i>n</i> -C ₇ H ₁₅ CO	H	H	D	411	
68 ²	H	Δ ⁹ -E <i>n</i> -C ₁₈ H ₃₅ CO	H	H	D	549	
69	COO- <i>i</i> -Pr	H	H	H	D	371	

TABLE 1

[illegible]

TABLE 1-A	
Cpd No.	Chemical Name
1	Acetic acid (2R,3S,4R,5R)-3,4-diacetoxy-5-(4-acetylamino-2-oxo-2H-pyrimidin-1-yl)-2-azido-tetrahydro-furan-2-ylmethyl ester
2	Propionic acid (2R,3S,4R,5R)-2-azido-5-(2-oxo-4-propionylamino-2H-pyrimidin-1-yl)-3,4-bis-propionyloxy-tetrahydro-furan-2-ylmethyl ester
3	Butyric acid (2R,3S,4R,5R)-2-azido-5-(4-butyrylamino-2-oxo-2H-pyrimidin-1-yl)-3,4-bis-butyryloxy-tetrahydro-furan-2-ylmethyl ester
4	3-Methyl-butyric acid (2R,3S,4R,5R)-2-azido-5-[4-(3-methyl-butyrylamino)-2-oxo-2H-pyrimidin-1-yl]-4-(3-methyl-butyryloxy)-2-(3-methyl-butyryloxymethyl)-tetrahydro-furan-3-yl ester
5	2,2-Dimethyl-propionic acid (2R,3S,4R,5R)-2-azido-5-[4-(2,2-dimethyl-propionylamino)-2-oxo-2H-pyrimidin-1-yl]-3,4-bis-(2,2-dimethyl-propionyloxy)-tetrahydro-furan-2-ylmethyl ester
6	Benzoic acid (2R,3S,4R,5R)-2-azido-3,4-dibenzoyloxy-5-(4-benzoylamino-2-oxo-2H-pyrimidin-1-yl)-tetrahydro-furan-2-ylmethyl ester
7	Phenoxy-acetic acid (2R,3S,4R,5R)-2-azido-5-[2-oxo-4-(2-phenoxy-acetylamino)-2H-pyrimidin-1-yl]-3,4-bis-(2-phenoxy-acetoxy)-tetrahydro-furan-2-ylmethyl ester
8	[1-((2R,3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-carbamic acid tert-butyl ester
9	N-[1-((2R,3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-acetamide
10	N-[1-((2R,3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-propionamide
11	N-[1-((2R,3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-butyramide
12	N-[1-((2R,3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-isobutyramide
13	N-[1-((2R,3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-2,2-dimethyl-propionamide
14	N-[1-((2R,3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-benzamide
15	N-[1-((2R,3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-2-phenyl-acetamide
16	N-[1-((2R,3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-2-phenoxy-acetamide
17	[1-((2R,3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-carbamic acid butyl ester
18	1-[(2R,3S,4R,5R)-5-(4-Amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethoxycarbonyl]-2-methyl-propyl-ammonium; chloride
19	1-[(2R,3S,4R,5R)-5-(4-Amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethoxycarbonyl]-2-phenyl-ethyl-ammonium; chloride
20	1-[(2R,3S,4R,5R)-5-(4-Amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethoxycarbonyl]-ethyl-ammonium; chloride
21	Benzoic acid (2R,3S,4R,5R)-5-(4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester
22	Acetic acid (2R,3S,4R,5R)-4-acetoxy-2-acetoxymethyl-5-(4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-tetrahydro-furan-3-yl ester
23	Propionic acid (2R,3S,4R,5R)-5-(4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-4-propionyloxy-2-propionyloxymethyl-tetrahydro-furan-3-yl ester
24	Isobutyric acid (2R,3S,4R,5R)-5-(4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-4-isobutyryloxy-2-isobutyryloxymethyl-tetrahydro-furan-3-yl ester
25	Benzoic acid (2R,3S,4R,5R)-3,4-dibenzoyloxy-5-(4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-tetrahydro-furan-2-ylmethyl ester
26	Benzoic acid (2R,3S,4R,5R)-3-(3-chloro-benzoyloxy)-4-benzoyloxy-5-(4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-tetrahydro-furan-2-ylmethyl ester
27	Benzoic acid (2R,3S,4R,5R)-3,4-dibenzoyloxy-2-azido-5-(4-butoxycarbonylamino-2-oxo-2H-pyrimidin-1-yl)-tetrahydro-furan-2-ylmethyl ester
28	Isobutyric acid (2R,3S,4R,5R)-2-azido-3,4-dihydroxy-5-[2-oxo-4-(2-phenoxy-acetylamino)-2H-pyrimidin-1-yl]-tetrahydro-furan-2-ylmethyl ester

TABLE 1-A

Cpd No.	Chemical Name
29	Butyric acid (3aS,4R,6R,6aR)-6-(4-amino-2-oxo-2H-pyrimidin-1-yl)-4-azido-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxol-4-ylmethyl ester
30	Benzoic acid (3aS,4R,6R,6aR)-6-(4-amino-2-oxo-2H-pyrimidin-1-yl)-4-azido-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxol-4-ylmethyl ester
31	2- <i>tert</i> -Butoxycarbonylamino-3-phenyl-propionic acid (3aS,4R,6R,6aR)-6-(4-amino-2-oxo-2H-pyrimidin-1-yl)-4-azido-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxol-4-ylmethyl ester
32	Isobutyric acid (3aS,4R,6R,6aR)-6-(4-amino-2-oxo-2H-pyrimidin-1-yl)-4-azido-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxol-4-ylmethyl ester
33	4-Amino-1-((3aR,4R,6R,6aS)-6-azido-6-hydroxymethyl-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxol-4-yl)-1H-pyrimidin-2-one
34	Isobutyric acid (3aS,4R,6R,6aR)-4-azido-6-(4-isobutyrylamino-2-oxo-2H-pyrimidin-1-yl)-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxol-4-ylmethyl ester
35	Acetic acid (3aS,4R,6R,6aR)-6-(4-acetylamino-2-oxo-2H-pyrimidin-1-yl)-4-azido-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxol-4-ylmethyl ester
36	Benzoic acid (3aS,4R,6R,6aR)-4-azido-6-(4-benzoylamino-2-oxo-2H-pyrimidin-1-yl)-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxol-4-ylmethyl ester
37	Carbonic acid (2R,3R,4S,5R)-5-azido-2-(4-butoxycarbonylamino-2-oxo-2H-pyrimidin-1-yl)-4-butoxycarbonyloxy-5-butoxycarbonyloxy-methyl-tetrahydro-furan-3-yl ester butyl ester
38 ¹	1-[1-((2R,3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-ylcarbonyl]-2-methyl-propyl-ammonium; chloride
39	Isobutyric acid (2R,3S,4R)-5-((S)-4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester
40 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-isobutyryloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
41 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-bis-isobutyryloxy-5-isobutyryloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
42	[(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-carbamic acid octyl ester
43 ²	(S)-1-((3R,4S,5R)-5-Azido-5-benzoyloxymethyl-3,4-dihydroxy-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
44	[(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-carbamic acid heptyl ester
45 ²	(S)-1-[(3R,4S,5R)-5-Azido-3,4-bis-(3-methyl-butyryloxy)-5-(3-methyl-butyryloxymethyl)-tetrahydro-furan-2-yl]-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
46 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-bis-pentanoyloxy-5-pentanoyloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
47	[(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-carbamic acid hexyl ester
48	Isobutyric acid (2R,3S,4R,5R)-5-(4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-2-hydroxymethyl-4-isobutyryloxy-tetrahydro-furan-3-yl ester
49 ²	1-[(2R,3S,4R)-5-((S)-4-Amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethoxycarbonyl]-2-methyl-butyl-ammonium; chloride
50 ²	(S)-1-[(2R,3S,4R)-5-azido-3,4-bis-(2,2-dimethyl-propionyloxy)-5-(2,2-dimethyl-propionyloxymethyl)-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
51	Carbonic acid (2R,3S,4R)-5-((S)-4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester propyl ester
52	Carbonic acid (2R,3S,4R)-5-((S)-4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester pentyl ester
53	Carbonic acid (2R,3S,4R)-5-((S)-4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester isobutyl ester
54 ²	(S)-1-((3R,4S,5R)-3,4-Diacetoxy-5-acetoxymethyl-5-azido-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
55	Isobutyric acid (2R,3S,4R)-2-azido-5-((S)-4-isobutyrylamino-2-oxo-2H-pyrimidin-1-yl)-4-isobutyryloxy-2-isobutyryloxymethyl-tetrahydro-furan-3-yl ester
56 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-bis-isobutyryloxy-5-isobutyryloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
57	[(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-carbamic acid methyl ester

TABLE 1-A

Cpd No.	Chemical Name
58	[(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-carbamic acid ethyl ester
59 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-bis-propionyloxy-5-propionyloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
60	Propionic acid (2R,3S,4R)-2-azido-5-((S)-4-benzoylamino-2-oxo-2H-pyrimidin-1-yl)-4-propionyloxy-2-propionyloxymethyl-tetrahydro-furan-3-yl ester
61 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-propoxycarbonyloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
62 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-bis-benzoyloxy-5-benzoyloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
63 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-pentyloxycarbonyloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
64 ³	(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hexyloxycarbonyloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; trifluoro-acetate
65 ³	(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-heptyloxycarbonyloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; trifluoro-acetate
66 ³	(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-octyloxycarbonyloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; trifluoro-acetate
67 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-octanoyloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
68 ²	(S)-1-[(3R,4S,5R)-5-Azido-3,4-dihydroxy-5-((E)-octadec-9-enoyloxymethyl)-tetrahydro-furan-2-yl]-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
69	[(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl]-carbamic acid isopropyl ester
70 ²	(S)-1-((3R,4S,5R)-5-Azido-5-heptanoyloxymethyl-3,4-dihydroxy-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
71 ²	(S)-1-((3R,4S,5R)-5-Azido-5-dodecanoyloxymethyl-3,4-dihydroxy-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
72	Undecanoic acid (2R,3S,4R)-5-((S)-4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester
73 ²	(S)-1-((3R,4S,5R)-5-Azido-5-decanoyloxymethyl-3,4-dihydroxy-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
74	Tetradecanoic acid (2R,3S,4R)-5-((S)-4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester
75	Nonanoic acid ((2R,3S,4R)-5-((S)-4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester
76	Heptanoic acid ((2R,3S,4R)-5-((S)-4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester
77	Hexadecanoic acid (2R,3S,4R)-5-((S)-4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester
78	Z-Octadec-9-enoic acid (2R,3S,4R)-5-((S)-4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester
79	3-Methyl-butyric acid (2R,3S,4R)-2-azido-5-((S)-4-butoxycarbonylamino-2-oxo-2H-pyrimidin-1-yl)-3,4-bis-(3-methyl-butyryloxy)-tetrahydro-furan-2-ylmethyl ester
80	Propionic acid (2R,3S,4R)-2-azido-5-((S)-4-butoxycarbonylamino-2-oxo-2H-pyrimidin-1-yl)-3,4-bis-propionyloxy-tetrahydro-furan-2-ylmethyl ester
81 ³	(S)-1-((3R,4S,5R)-5-Azido-3,4-bis-hexanoyloxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; trifluoro-acetate
82 ³	(S)-1-((3R,4S,5R)-5-azido-5-hydroxymethyl-3,4-bis-pentanoyloxy-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; trifluoro-acetate
83 ²	(S)-1-((3R,4S,5R)-5-Azido-5-hydroxymethyl-3,4-bis-pentanoyloxy-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
84 ²	(S)-1-[(3R,4S,5R)-5-Azido-3,4-dihydroxy-5-(4-methyl-benzoyloxymethyl)-tetrahydro-furan-2-yl]-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
85 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-tridecanoyloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
86 ³	(S)-1-((3R,4S,5R)-5-Azido-5-decyloxycarbonyloxymethyl-3,4-dihydroxy-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; trifluoro-acetate

TABLE 1-A	
Cpd No.	Chemical Name
87	Dodecanoic acid (2R,3S,4R)-5-((S)-4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester
88 ³	(S)-1-((3R,4S,5R)-5-azido-3,4-bis-butyryloxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; trifluoro-acetate
89 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-bis-butyryloxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
90 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-nonyloxycarbonyloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
91 ²	(S)-1-((3R,4S,5R)-5-Azido-5-hydroxymethyl-3,4-bis-isobutyryloxy-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
92 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-bis-benzoyloxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
93 ²	(S)-1-((3R,4S,5R)-5-Azido-5-hydroxymethyl-3,4-bis-propionyloxy-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
94 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-bis-butyryloxy-5-butyryloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
95 ³	(S)-1-((3R,4S,5R)-5-azido-3,4-bis-propionyloxy-5-propionyloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; methanesulfonate
96	Carbonic acid (2R,3S,4R)-2-azido-3,4-dihydroxy-5-((S)-2-oxo-4-pentyloxycarbonylamino-2H-pyrimidin-1-yl)-tetrahydro-furan-2-ylmethyl ester pentyl ester
97	Carbonic acid (2R,3S,4R)-2-azido-5-((S)-4-hexyloxycarbonylamino-2-oxo-2H-pyrimidin-1-yl)-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester hexyl ester
98 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-octylcarbonyloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride
99	Benzoic acid (2R,3S,4R)-2-azido-5-((S)-4-hexanoylamino-2-oxo-2H-pyrimidin-1-yl)-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester
100	Benzoic acid (2R,3S,4R)-2-azido-5-((S)-4-heptanoylamino-2-oxo-2H-pyrimidin-1-yl)-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester
101 ³	(S)-1-((3R,4S,5R)-5-azido-5-hydroxymethyl-3,4-bis-pentanoyloxy-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; methanesulfonate
102 ²	(S)-1-((3R,4S,5R)-5-azido-3,4-bis-hexanoyloxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; methanesulfonate
103 ²	(S)-1-((3R,4S,5R)-5-Azido-3,4-dihydroxy-5-pentanoyloxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl-ammonium; chloride

The compounds of formula I may be prepared by various methods known in the art of organic chemistry in general and nucleoside analogue synthesis in particular. Specific methodologies to prepare compounds of the present invention are illustrated in Examples

- 5 1-6. The starting materials for the syntheses are either readily available from commercial sources or are known or may themselves be prepared by techniques known in the art. General reviews of the preparation of nucleoside analogues are included in the following publications which are incorporated herein in their entirety:

A. M Michelson "The Chemistry of Nucleosides and Nucleotides", Academic Press, New
10 York 1963.

L. Goodman "Basic Principles in Nucleic Acid Chemistry" Ed P O P Ts'O, Academic Press, New York 1974, Vol. 1, chapter 2.

"Synthetic Procedures in Nucleic acid Chemistry" Ed W W Zorbach and R S Tipson, Wiley, New York, 1973, Vol. 1 and 2.

H.Vorbrüggen and C. Ruh-Pohlenz (eds) "Handbook of Nucleoside Synthesis" Wiley, New York, 2001.

- 5 The compounds of the present invention are prepared by acylation of a suitable nucleoside compound. Acylation of alcohols (J. March, *Advanced Organic Chemistry* John Wiley & Sons, New York 1992 392-398; J. Mulzer *Synthesis of Esters, Activated Esters & Lactones in Comprehensive Organic Synthesis*, E. Winterfeldt, ed., vol. 6, Pergamon Press, Oxford 1991, pp.324-340) and amines (J. March, *supra* pp.417-425; H. G. Benz, 10 *Synthesis of Amides and Related Compounds in Comprehensive Organic Synthesis*, E. Winterfeldt, ed., vol. 6, Pergamon Press, Oxford 1991 pp. 381-411) can be accomplished with a variety of acylating agents including acid chlorides and acid anhydrides. These references are incorporated herein in their entirety. Other methods for activation of a carboxylic acid have been developed and can be utilized to prepare the prodrugs 15 described herein. The extent and pattern of acylation is controlled by use of suitable protecting groups or by the delaying the introduction of the basic amine into the pyrimidine base.

Tetraacyl nucleosides are readily prepared by acylating 1-((2R,3R,4S,5R)-5-azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-pyrimidin-4-yl- 20 ammonium; hydrogen sulfate (II) with at least 4 equivalents of acylating agent (Method A; Example 1).

Amines typically are more reactive toward acylating agents than are hydroxyl groups. Nonetheless to insure selective acylation of the amine substituent the hydroxyl groups are protected prior to acylation (Method B; Example 2). Trimethylsilyl ethers are useful 25 protecting groups for this transformation. More detailed information regarding protection and deprotection of alcohols and alternative protecting groups can be found in T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, 3rd edition, John Wiley & Sons, New York, 1999, and Harrison and Harrison et al., *Compendium of Synthetic Organic Methods*, Vols. 1-8 John Wiley and Sons, 1971-1996. The above 30 references are incorporated herein by reference in their entirety.

Selective acylation of the three hydroxy substituents is accomplished by carrying out the acylation on the corresponding uridine nucleoside II, which lacks the reactive amine substituent on the heteroaromatic ring, and subsequently converting the uridine to a cytidine. The acylated uridine can be converted to cytidine by utilizing the method

described by A. D. Borthwick *et al.*, (*J. Med. Chem.* 1990, 33(1):179; see also K. J. Divakar and C. B. Reese *J. Chem Soc., Perkin Trans. I* 1982 1171-1176 and Maag *et al. J. Med. Chem.* 1992 35:1440-1451). The invention further provides a method for the selective cleavage of the N-acyl moiety from a N acylated nucleoside by contacting the N-acyl
5 compound with zinc bromide in a protic solvent.

Selective acylation of the 5-hydroxy group and the amine was accomplished by protection of the *vicinal* 2',3'-hydroxy groups of a carbohydrate before acylation of the primary alcohol (Method E; Example 5). Protecting groups for *vicinal* diols often convert the diol into a dioxolane or dioxane ring (see Greene *supra*; Harrison and Harrison *supra*). Most
10 commonly these protecting groups include aldehydes and ketones which readily form dioxolanes. Ketones which have found particular utility as diol protecting groups include acetone and C₅₋₇ cycloalkanones. Cleavage of the dioxolane or dioxane to regenerate the diol is generally accomplished with aqueous acid and an organic cosolvent. Benzaldehyde readily forms acetals with *vic* diols which can be deprotected by hydrogenolysis or acidic
15 hydrolysis. Methoxy substitution on the benzaldehyde increases the rate of acidic hydrolysis and also permits cleavage of the dioxolane under oxidative conditions, e.g. Ce(NH₄)₂(NO₃)₆. Nitrobenzaldehydes afford dioxolanes which can be photochemically cleaved. Cyclic orthoesters, e.g. ethoxymethylene acetal have been utilized as diol protecting groups. These compounds can be cleaved under mild acidic conditions;
20 however the initial product is an ester which must be hydrolyzed to regenerate the diol. The cyclic analog 2-oxacyclopentylidene orthoester affords the diol directly upon acid hydrolysis. Cyclic carbonates and cyclic boronates also have found some utility as diol protecting groups. Any of these diol protecting groups can be adapted to the present process.

25 5'-Monoacyl derivatives are accessible by protection of the 2',3' *vicinal* diol of a uridine derivative. Acylation of the remaining reactive hydroxyl is followed by conversion of the uridine base to the corresponding cytidine base as described above and deprotection of the *vicinal* diol (Method D; Example 4). Alternatively the N-acyl group of an N,1'-diacyl cytidine compound in which the 3' and 4' alcohols are protected can be selectively cleaved
30 with zinc bromide to produce the protected monoacyl compound which can be further deprotected (R. Kierzek *et al. Tetrahedron Lett.* 1981 22(38):3762-64).

Compounds of the present invention may be administered either alone (*i.e.*, monotherapy) or in combination with other therapeutic agents (e.g., "combination therapy"). Combination therapy can consist of an HCV polymerase inhibitor and
35 immune modulators which stimulate natural immune responses to the virus and viral infected cells such as interferons, chemically modified interferons, interleukin, tumor necrosis factor or colony stimulating factors. Compounds of the present invention also

can be combined with other antiviral compounds with a similar or complementary mode of action. Potential targets for antiviral drugs have been reviewed. (see e.g., E. DeClercq, *Strategies in the Design of Antiviral Drugs*, *Nature Rev Drug Discov.* 2002 1(1):13-25; M. T. Reding, *Recent Developments in hepatitis C antiviral research 1999-2000*, *Exp Opin.*

- 5 *Therap. Pat.* 2000, 10(8):1201-1220) Antiviral compounds including, but not limited to, HCV protease inhibitors, other HCV polymerase inhibitors, HCV helicase inhibitors, HCV primase inhibitors, HCV integrase inhibitors or HCV fusion inhibitors all could be useful in combination with compounds of the present invention.

- The compounds of the present invention may be formulated in a wide variety of oral
10 administration dosage forms and carriers. While nucleoside derivatives of the present invention are optimized for delivery across the gastrointestinal mucosa, these compounds can be efficacious when administered by other routes of administration. The pharmaceutically acceptable carriers may be either solid or liquid. Oral administration can be in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules,
15 solutions, emulsions, syrups, or suspensions. Compounds of the present invention are efficacious when administered by other routes of administration including continuous (intravenous drip) topical parenteral, intramuscular, intravenous, subcutaneous, transdermal (which may include a penetration enhancement agent), buccal, nasal, inhalation and suppository administration, among other routes of administration. The
20 preferred manner of administration is generally oral using a convenient daily dosing regimen which can be adjusted according to the degree of affliction and the patient's response to the active ingredient.

- A compound or compounds of the present invention, as well as their pharmaceutically useable salts, together with one or more conventional excipients, carriers, or diluents,
25 may be placed into the form of pharmaceutical compositions and unit dosages. The pharmaceutical compositions and unit dosage forms may be comprised of conventional ingredients in conventional proportions, with or without additional active compounds or principles, and the unit dosage forms may contain any suitable effective amount of the active ingredient commensurate with the intended daily dosage range to be employed.
30 The pharmaceutical compositions may be employed as solids, such as tablets or filled capsules, semisolids, powders, sustained release formulations, or liquids such as solutions, suspensions, emulsions, elixirs, or filled capsules for oral use; or in the form of suppositories for rectal or vaginal administration; or in the form of sterile injectable solutions for parenteral use. A typical preparation will contain from about 5% to about
35 95% active compound or compounds (w/w).

The term "excipient" as used herein refers to a compound that is useful in preparing a pharmaceutical composition, generally safe, non-toxic and neither biologically nor otherwise undesirable, and includes excipients that are acceptable for veterinary use as well as human pharmaceutical use. The term "excipient" as used herein includes both one
5 and more than one such excipient.

Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier may be one or more substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material. In powders, the
10 carrier generally is a finely divided solid which is a mixture with the finely divided active component. In tablets, the active component generally is mixed with the carrier having the necessary binding capacity in suitable proportions and compacted in the shape and size desired. Suitable carriers include but are not limited to magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth,
15 methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. Solid form preparations include solutions, suspensions, and emulsions, and may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

The term "preparation" or "dosage form" as used herein is intended to include the
20 formulation of the active compound with encapsulating material as carrier, providing a capsule in which the active component, with or without carriers, is surrounded by a carrier, which is in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges may be as solid forms suitable for oral administration.

25 Liquid formulations also are suitable for oral administration include liquid formulation including emulsions, syrups, elixirs, aqueous solutions, aqueous suspensions are also suitable forms for oral administration. These include solid form preparations which are intended to be converted to liquid form preparations shortly before use. Emulsions may be prepared in solutions, for example, in aqueous propylene glycol solutions or may
30 contain emulsifying agents such as lecithin, sorbitan monooleate, or acacia. Aqueous solutions can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing, and thickening agents. Aqueous suspensions can be prepared by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium
35 carboxymethylcellulose, and other well known suspending agents.

The compounds of the present invention may be formulated for parenteral administration (*e.g.*, by injection, for example bolus injection or continuous infusion) and may be presented in unit dose form in ampoules, pre-filled syringes, small volume infusion or in multi-dose containers with an added preservative. The compositions may
5 take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, for example solutions in aqueous polyethylene glycol. Examples of oily or nonaqueous carriers, diluents, solvents or vehicles include propylene glycol, polyethylene glycol, vegetable oils (*e.g.*, olive oil), and injectable organic esters (*e.g.*, ethyl oleate), and may contain formulatory agents such as preserving, wetting, emulsifying or suspending,
10 stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form, obtained by aseptic isolation of sterile solid or by lyophilisation from solution for constitution before use with a suitable vehicle, *e.g.*, sterile, pyrogen-free water.

The compounds of the present invention may be formulated for topical administration to the epidermis as ointments, creams or lotions, or as a transdermal patch. Ointments and
15 creams may, for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Lotions may be formulated with an aqueous or oily base and will in general also containing one or more emulsifying agents, stabilizing agents, dispersing agents, suspending agents, thickening agents, or coloring agents. Formulations suitable for topical administration in the mouth include lozenges
20 comprising active agents in a flavored base, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert base such as gelatin and glycerin or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

The compounds of the present invention may be formulated for administration as
25 suppositories. A low melting wax, such as a mixture of fatty acid glycerides or cocoa butter is first melted and the active component is dispersed homogeneously, for example, by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and to solidify.

The compounds of the present invention may be formulated for vaginal administration.
30 Pessaries, tampons, creams, gels, pastes, foams or sprays containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

The compounds of the present invention may be formulated for nasal administration. The solutions or suspensions are applied directly to the nasal cavity by conventional means, for example, with a dropper, pipette or spray. The formulations may be provided

in a single or multidose form. In the latter case of a dropper or pipette, this may be achieved by the patient administering an appropriate, predetermined volume of the solution or suspension. In the case of a spray, this may be achieved for example by means of a metering atomizing spray pump.

- 5 The compounds of the present invention may be formulated for aerosol administration, particularly to the respiratory tract and including intranasal administration. The compound will generally have a small particle size for example of the order of five (5) microns or less. Such a particle size may be obtained by means known in the art, for example by micronization. The active ingredient is provided in a pressurized pack with a
- 10 suitable propellant such as a chlorofluorocarbon (CFC), for example, dichlorodifluoromethane, trichlorofluoromethane, or dichlorotetrafluoroethane, or carbon dioxide or other suitable gas. The aerosol may conveniently also contain a surfactant such as lecithin. The dose of drug may be controlled by a metered valve. Alternatively the active ingredients may be provided in a form of a dry powder, for
- 15 example a powder mix of the compound in a suitable powder base such as lactose, starch, starch derivatives such as hydroxypropylmethyl cellulose and polyvinylpyrrolidone (PVP). The powder carrier will form a gel in the nasal cavity. The powder composition may be presented in unit dose form for example in capsules or cartridges of e.g., gelatin or blister packs from which the powder may be administered by means of an inhaler.
- 20 When desired, formulations can be prepared with enteric coatings adapted for sustained or controlled release administration of the active ingredient. For example, the compounds of the present invention can be formulated in transdermal or subcutaneous drug delivery devices. These delivery systems are advantageous when sustained release of the compound is necessary and when patient compliance with a treatment regimen is
- 25 crucial. Compounds in transdermal delivery systems are frequently attached to an skin-adhesive solid support. The compound of interest can also be combined with a penetration enhancer, e.g., Azone (1-dodecylazacycloheptan-2-one). Sustained release delivery systems are inserted subcutaneously into the subdermal layer by surgery or injection. The subdermal implants encapsulate the compound in a lipid soluble
- 30 membrane, e.g., silicone rubber, or a biodegradable polymer, e.g., polylactic acid.

Suitable formulations along with pharmaceutical carriers, diluents and excipients are described in *Remington: The Science and Practice of Pharmacy* 1995, edited by E. W. Martin, Mack Publishing Company, 19th edition, Easton, Pennsylvania. A skilled formulation scientist may modify the formulations within the teachings of the

35 specification to provide numerous formulations for a particular route of administration

without rendering the compositions of the present invention unstable or compromising their therapeutic activity.

The modification of the present compounds to render them more soluble in water or other vehicle, for example, may be easily accomplished by minor modifications (salt formulation, esterification, etc.), which are well within the ordinary skill in the art. It is also well within the ordinary skill of the art to modify the route of administration and dosage regimen of a particular compound in order to manage the pharmacokinetics of the present compounds for maximum beneficial effect in patients.

The phrase "therapeutically effective amount" as used herein means an amount required to reduce symptoms of the disease in an individual. The dose will be adjusted to the individual requirements in each particular case. That dosage can vary within wide limits depending upon numerous factors such as the severity of the disease to be treated, the age and general health condition of the patient, other medicaments with which the patient is being treated, the route and form of administration and the preferences and experience of the medical practitioner involved. For oral administration, a daily dosage of between about 0.01 and about 100 mg/kg body weight per day should be appropriate in monotherapy and/or in combination therapy. A preferred daily dosage is between about 0.1 and about 500 mg/kg body weight, more preferred 0.1 and about 100 mg/kg body weight and most preferred 1.0 and about 10 mg/kg body weight per day. Thus, for administration to a 70 kg person, the dosage range would be about 0.7 g to 7.0 g per day. The daily dosage can be administered as a single dosage or in divided dosages, typically between 1 and 5 dosages per day. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect for the individual patient is reached. One of ordinary skill in treating diseases described herein will be able, without undue experimentation and in reliance on personal knowledge, experience and the disclosures of this application, to ascertain a therapeutically effective amount of the compounds of the present invention for a given disease and patient.

The pharmaceutical preparations are preferably in unit dosage forms. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.


The pharmaceutical compositions in Example 8 are given to enable those skilled in the art to more clearly understand and to practice the present invention. They should not be considered as limiting the scope of the invention, but merely as being illustrative and representative thereof.

- 5 The following preparations and examples are given to enable those skilled in the art to more clearly understand and to practice the present invention. They should not be considered as limiting the scope of the invention, but merely as being illustrative and representative thereof.

- Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts,
10 temperatures), but allowance for some experimental error and deviation, including differences in calibration, rounding of numbers, and the like, is contemplated.

EXAMPLE 1

Method A: Preparation of N, 2',3',4'-Tetraacyl Nucleoside Derivatives




Acetic acid 3,4-diacetoxy-5-(4-acetylamino-2-oxo-2H-pyrimidin-1-yl)-2-azido-
tetrahydro-furan-2-ylmethyl ester

- 5 A stirred suspension of 4'-azidocytidine hemisulfate (0.30 g), dimethylaminopyridine (cat.), and N,N-diisopropylethylamine (3.69 mL) in methylene chloride (5 mL), under inert atmosphere, was cooled in an ice/water bath and treated dropwise with acetyl chloride (0.452 mL) and acetic anhydride (0.60 mL). The mixture was allowed to warm
 10 to ambient temperature and after 2 days was subjected, untreated, to flash chromatography (50% ethyl acetate in hexanes, then 75% ethyl acetate in hexanes, then 100% ethyl acetate, then 5% methanol in ethyl acetate) to afford 0.335 g of the solid product (compound 1; M+H=453).

EXAMPLE 2

- 15 **Method B: Preparation of N-acyl Nucleoside Derivatives**
[1-(5-Azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2-oxo-1,2-dihydro-
pyrimidin-4-yl]-carbamic acid butyl ester




- 34 -

A stirred suspension of 4'-azidocytidine hemisulfate (0.50g) in pyridine (8 mL), under inert atmosphere, was cooled in an ice/water bath and treated with trimethylsilyl chloride (1 mL). The mixture was allowed to warm to ambient temperature and after 1h was treated with butyl chloroformate (0.2 mL). After stirring another 2h at that temperature, the reaction was cooled in an ice/water bath and treated with 5 mL aqueous ammonium bicarbonate. The organics were extracted twice with methylene chloride, dried over magnesium sulfate, and filtered. To the filtrate was added tetra-n-butyl ammonium fluoride (0.25 mL, 1 M in tetrahydrofuran). The reaction was stirred at ambient temperature for three days. After solvent removal, the residue was subjected to flash chromatography (25% hexanes in ethyl acetate, then ethyl acetate, then 6.5% methanol in ethyl acetate) to afford 400 mg of the product as a solid (compound 17; mp 86.5-94 °C).

EXAMPLE 3

Method C: Preparation of 2', 3', 5'-Triacyl Nucleoside Derivatives

Acetic acid 3,4-diacetoxy-5-(4-amino-2-oxo-2H-pyrimidin-1-yl)-tetrahydro-furan-2-ylmethyl ester (22)

Step 1

To a stirred solution containing 0.330 g (1.15 mmol) 4'-azidouridine, 2 mL pyridine and 2 mL acetic anhydride was added 0.010 g (0.08 mmol) of 4-dimethylaminopyridine. After

12 h, the reaction mixture was evaporated to dryness under reduced pressure. The residue was dissolved in dichloromethane and washed with saturated aqueous sodium hydrogen carbonate solution, dried over magnesium sulfate and evaporated to dryness to give 0.42 g (88%) of 2',3',5'-tri-acetoxy-4'-azidouridine (IIIa: $R^5 = CH_3$).

5 Step 2

$POCl_3$ (0.31 mL; 3.304 mmol) was added to a stirred mixture containing 0.340 g (0.826 mmol) of the uridine, 0.913 g (13.22 mmol) of 1,2,4-triazole and 2.30 mL (16.52 mmol) of triethylamine in 20 mL acetonitrile cooled to 5 °C. The reaction mixture was allowed to warm to room temperature. After 12 h, the reaction mixture was evaporated to dryness
10 under reduced pressure. The residue was diluted with dichloromethane, washed with saturated aqueous sodium hydrogen carbonate solution, dried over magnesium sulfate and evaporated to dryness to give 0.300 g (78%) of (IIc: $R^5 = CH_3$).

Step 3


To a sealed flask containing 0.066 g (1.254 mmol) ammonium chloride and 0.070 g
15 (1.254 mmol) of potassium hydroxide was added 10 mL acetonitrile, 20 mL water, 0.190 mL (1.278 mmol) of triethylamine followed by a solution containing 0.290 g (0.627 mmol) of the triazole in 10 mL acetonitrile. After 12 h, the mixture was evaporated to dryness under reduced pressure. The residue was dissolved in ethyl acetate and washed with water, dried over magnesium sulfate and evaporated to dryness under reduced
20 pressure. Chromatography (10% methanol in dichloromethane) provided 0.060 g (23%) of the cytidine. (IIId; $R^5 = CH_3$; compound 22; $MH^+ = 411$; $MNa^+ = 433$).

Proceeding in similar fashion with the appropriate acylating agent there was obtained isobutyric acid 5-(4-amino-2-oxo-2H-pyrimidin-1-yl)-3,4-bis-isobutyryloxy-tetrahydro-furan-2-ylmethyl ester (compound 24; $MH^+ = 495$; $MNa^+ = 517$) and benzoic acid 5-(4-
25 amino-2-oxo-2H-pyrimidin-1-yl)-3,4-di-benzoxo-tetrahydro-furan-2-ylmethyl ester (compound 25).

EXAMPLE 4

Method D: Preparation of 5'-Acyl Nucleoside Derivatives

- 2-Amino-3-phenyl-propionic acid 5-(4-amino-2-oxo-2H-pyrimidin-1-yl)-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester hydrochloride (Ie: R = CH(NH₃⁺)CH₂C₆H₅ Cl; compound 19)

Step 1

- 1-(6-Hydroxymethyl-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxol-4-yl)-1H-pyrimidine-2,4-dione (IIIb; compound 33)

A mixture containing 3.0 g (10.5 mmol) of 4'-azidouridine, 0.05 g (0.26 mmol) *p*-toluenesulfonic acid monohydrate and 6 mL (48.8 mmol) 2,2-dimethoxypropane in 20 mL acetone was stirred at room temperature for 12 h. The reaction mixture was diluted with ethyl acetate, washed with saturated aqueous sodium hydrogen carbonate solution, dried over magnesium sulfate and evaporated to dryness under reduced pressure to give the desired product 2.20 g (64%) of as a white solid.

Step 2

- 2-*tert*-Butoxycarbonylamino-3-phenyl-propionic acid 6-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxol-4-ylmethyl ester (IIIb: R' = CH(NH-boc)CH₂C₆H₅)

- 37 -

To a stirred solution containing 1.00 g (3.07 mmol) of 4'-azido-2',3'-O-iso-propyl-uridine, 1.63 g (6.14 mmol) Boc-L-phenylalanine and 1.18 g (6.14 mmol) 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride in 20 mL *N,N*-dimethylformamide was added 0.375 g (3.07 mmol) of 4-dimethylaminopyridine. The
 5 resulting solution was left to stir under an atmosphere of nitrogen and at room temperature. After 12 h, the reaction mixture was evaporated to dryness under reduced pressure. Chromatography (0 to 100% ethyl acetate in hexanes) of the crude residue gave 1.01 g (57%) of the desired product as a white foam ($MH^+ = 573$; $MNa^+ = 595$).

Step 3

10 2-tert-Butoxycarbonylamino-3-phenyl-propionic acid 6-(4-amino-2-oxo-2H-pyrimidin-1-yl)-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxol-4-ylmethyl ester (IIe: R' = CH(NHboc)CH₂C₆H₅)

POCl₃ (0.651 mL; 6.98 mmol) was added to a stirred mixture containing 1.00 g (1.746 mmol) of the uridine, 1.93 g (27.94 mmol) of 1,2,4-triazole and 4.86 mL (34.93
 15 mmol) of triethylamine in 50 mL acetonitrile cooled to 5 °C. The reaction mixture was allowed to warm to room temperature. After 12 h, the reaction mixture was evaporated to dryness under reduced pressure. The residue was diluted with dichloromethane, washed with saturated aqueous sodium hydrogen carbonate solution, dried over magnesium sulfate and evaporated to dryness to give 1.0 g (92%) of the triazole.

20 To a sealed flask containing 0.171 g (3.207 mmol) ammonium chloride and 0.180 g (3.207 mmol) of potassium hydroxide was added 10 mL acetonitrile, 20 mL water, 0.446 mL (3.207 mmol) of triethylamine followed by a solution containing 1.00 g (1.603 mmol) of the phenylalanineuridine in 10 mL acetonitrile. After 12 h, the mixture was evaporated to dryness under reduced pressure. The residue was dissolved in ethyl acetate and washed
 25 with water, dried over magnesium sulfate and evaporated to dryness under reduced pressure. Chromatography (10% methanol in dichloromethane) provided 0.48 g (52%) of the cytidine. ($MH^+ = 572$; $MNa^+ = 594$).

Step 4:

2-Amino-3-phenyl-propionic acid 5-(4-amino-2-oxo-2H-pyrimidin-1-yl)-3,4-
 30 dihydroxy-tetrahydro-furan-2-ylmethyl ester hydrochloride (Ie: R' = CH(NH₃⁺)CH₂C₆H₅ Cl⁻)

To a solution containing 0.23 g (0.402 mmol) of phenylalanine cytidine (IIe: R' = CH(NHBoc)CH₂Ph) in 10 mL of methanol was added 0.079 mL (0.804 mmol) concentrated hydrogen chloride solution. After 12 h, the reaction mixture was evaporated
 35 to dryness. The residue was dissolved in water and washed with ethyl acetate and

- 38 -

evaporated to dryness under reduced pressure to give 0.160g (94%) cytidine product (compound 19).


In similar fashion substituting boc-L-valine and boc-L-alanine there was obtained respectively, 2-amino-3-methyl-butyrilic acid 5-(4-amino-2-oxo-2*H*-pyrimidin-1-yl)-3,4-
5 dihydroxy-tetrahydro-furan-2-ylmethyl ester hydrochloride (compound 18; $MH^+ = 384$; $MNa^+ = 406$) and 2-amino-propionic acid 5-(4-amino-2-oxo-2*H*-pyrimidin-1-yl)-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester hydrochloride (compound 20; $MH^+ = 356$; $MNa^+ = 378$).

In similar manner utilizing benzoic anhydride there was obtained benzoic acid 5-(4-
10 amino-2-oxo-2*H*-pyrimidin-1-yl)-3,4-dihydroxy-tetrahydro-furan-2-ylmethyl ester (compound 14; $MH^+ = 389$; $MNa^+ = 411$).

EXAMPLE 5

Method E

Isobutyric acid 2-azido-3,4-dihydroxy-5-(2-oxo-4-phenylacetyl-amino-2H-pyrimidin-1-yl)-tetrahydro-furan-2-ylmethyl ester



5

Step 1

A solution of 4-amino-1-(6-azido-6-hydroxymethyl-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxol-4-yl)-1H-pyrimidin-2-one (II, 0.14g), 4-nitrophenyl phenylacetate (0.12 g) and 1-hydroxybenzotriazole (0.06 g) in DMF (15 mL) was stirred at room temperature overnight. Water (15 mL) was added and the mixture was twice extracted with ethyl acetate. The combined ethyl acetate extracts were washed with water and brine and dried over anhydrous magnesium sulfate. The solution was concentrated *in vacuo* and purified by flash column chromatography which yielded IIh (0.18g) as a colorless oil.

15 Step 2

IIh from the previous step (0.18 g), triethylamine (0.07 ml) and dimethylaminopyridine (0.01 g) were dissolved in THF (15 ml) and stirred at room temperature under nitrogen atmosphere. Isobutyryl chloride (0.043 mL) was added slowly and the reaction was stirred at rt for 6 hours. Water (15 mL) was added and the mixture was extracted twice

with ethyl acetate. The combined extracts were dried over anhydrous magnesium sulfate, filtered, and the solution concentrated *in vacuo*. The product (IIi; 0.18 g) was separated by a flash column chromatography as an oil.


Step 3

- 5 IIi (0.18 g) was dissolved in acetic acid (60%) and stirred at 100 °C overnight. The reaction was cooled to rt and the solvent evaporated under reduced pressure. The product was purified by preparatory thin layer chromatography. The crude product (69 mg) was dissolved in THF (10 ml) and treated at room temperature with 4-nitrophenyl phenylacetate (55 mg) and 1-hydroxybenzotriazole (26 mg). The reaction mixture was
 10 stirred at rt overnight. The solvent was evaporated and the residue was re-purified by preparatory thin layer chromatography to afford IIj (17 mg) as a colorless oil.

EXAMPLE 6

Method F

- 15 Isobutyric acid (2R,3S,4R,5R)-5-(4-amino-2-oxo-2H-pyrimidin-1-yl)-2-azido-2-hydroxymethyl-4-isobutyryloxy-tetrahydro-furan-3-yl ester



Step 1

- A flask was charged with II (0.610 g; 1.75 mmol), dimethyl-*tert*-butylsilane (0.580 g; 3.84 mmol), imidazole (0.262 g; 3.842 mmol) and DMF (12 mL) and the resulting
 20 homogenous solution was stirred overnight at rt. Mass spectroscopy indicated the crude

- 41 -

product was a mixture of mono- and disilylated product. The solvent was removed *in vacuo* and the residue was dissolved in pyridine (12 mL) and CH₂Cl₂ (12 mL) and a small catalytic quantity of DMAP and isobutyric anhydride (0.96 mL; 5.76 mmol) was added and the resulting mixture was stirred overnight at rt. The reaction mixture was extracted with a mixture of water and saturated NaHCO₃ and the aqueous layer back-extracted CH₂Cl₂. The organic layers were combined and extracted with 1N HCl and water, dried (MgSO₄), filtered and evaporated to yield 1.3 g of crude silylated nucleosides as a yellow oil.

Step 2

The crude product (1.3 g) from the previous step was dissolved in THF (20 mL) and tetrabutyl ammonium fluoride (0.5 mL; TBAF; 1.0M solution in THF) was added and the reaction mixture was allowed to stir overnight. After 16 h, another 0.5 mL aliquot of TBAF was added and stirring continued for another 4 h and the solvent was evaporated. The residue was partitioned between CH₂Cl₂ and a mixture of H₂O and saturated NaHCO₃. The aqueous extract was washed with CH₂Cl₂ and the combined organic extracts were washed with H₂O and aqueous NaCl, dried, filtered and evaporated. The crude product was purified by flash chromatography and eluted with a gradient (CH₂Cl₂ → 4% MeOH/CH₂Cl₂) to yield a colorless white solid N, 2',3'-tri-isobutyrate (III_k; 140 mg)

Step 3

The tri-isobutyrate (140 mg; 0.283 mmol) was dissolve in 2.5 mL CH₂Cl₂ and 0.8 mL of MeOH. To the resulting solution was added ZnBr₂ (6 mg; 0.283 mmol) and the resulting solution was stirred at 65° C overnight. The solvent was removed *in vacuo* to yield 150 mg of a yellow foam which was purified by flash chromatography on silica gel and eluted with a gradient (CH₂Cl₂ → 75% MeOH/CH₂Cl₂) to yield II_m (R = *i*-Pr) as a white solid (0.120g; 98 % theory).

EXAMPLE 7

Plasma Pharmacokinetics

Pharmacokinetic procedures were used to determine plasma levels of 4-amino-1-((2R,3R,4S,5R)-5-azido-3,4-dihydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-1H-pyrimidin-2-one (II) after administration of a single oral 5 mg/kg dose of a prodrug of II. The formulation is a solution / suspension containing 0.0176 mmol of prodrug in 5 mL of an appropriate vehicle.

Three unfasted male *Cynomolgus* monkeys (6-9 kg) were fitted with a saphenous or

brachial catheter to facilitate blood draw. Free access to food and water will be allowed at all times during the study. On the day of the study a predose blood sample (2-3 mL) was taken from each monkey. The monkeys were dosed with 1 mL/kg of the dose solution by oral gavage. At each of the following time points (0.25, 0.5, 1, 3, 5, 7, 10, 24, 32, and 48
 5 hour) after dosing, approximately 0.5 mL of blood will be collected into lithium heparin-coated tubes. Blood was centrifuged to obtain plasma which was frozen until analysis.

The concentration of compound I, wherein R¹-R⁴ are H in each plasma sample was determined by an LC-MS assay. Standard curves were prepared in blank monkey plasma. The AUC represents the area under a plot of concentration vs time total which describes
 10 the concentration of the drug in systemic circulation as a function of time (L. Z. Benet, D. L. Kroetz and L. B. Sheiner *Pharmacokinetics in Goodman & Gilman's The Pharmacological Basis of Therapeutics*, J.G. Hardman & L.E. Limbird Eds., 9th Edition, McGraw Hill, New York, p 17-23). C_{max} is the peak concentration which is found.

Cpd No.	AUC _{0-24h} (µg·h/mL)	AUC fold increase	C _{max} (µg/mL)	C _{max} fold increase
II	2.85	1.00	0.312	1.00
22	3.34	1.22	0.378	1.21
49	2.38	0.87	0.217	0.70
18	2.37	0.86	0.284	0.91
50	1.64	0.60	0.171	0.55
23	9.50	3.47	2.09	6.70
51	0.89	0.32	0.081	0.26
52	1.40	0.51	0.308	0.99
42	0.65	0.24	0.074	0.24
53	0.58	0.21	0.058	0.19
47	0.83	0.30	0.103	0.33
46	6.99	2.55	1.11	3.56
44	5.21	1.90	0.827	2.65
24	10.5	3.83	1.42	4.55

15

EXAMPLE 8

Renilla luciferase assay

This assay measures the ability of the compounds of formula I to inhibit HCV RNA replication, and therefore their potential utility for the treatment of HCV infections. The assay utilizes a reporter as a simple readout for intracellular HCV replicon RNA level. The
 20 *Renilla* luciferase gene was introduced into the first open reading frame of a replicon construct NK5.1 (Krieger *et al.*, *J. Virol.* 75:4614), immediately after the internal ribosome entry site (IRES) sequence, and fused with the neomycin phosphotransferase (NPTII)

- gene via a self-cleavage peptide 2A from foot and mouth disease virus (Ryan & Drew, EMBO Vol 13:928-933). After *in vitro* transcription the RNA was electroporated into human hepatoma Huh7 cells, and G418-resistant colonies were isolated and expanded. Stably selected cell line 2209-23 contain replicative HCV subgenomic RNA, and the
- 5 activity of *Renilla* luciferase expressed by the replicon reflects its RNA level in the cells. The assay was carried out in duplicate plates, one in opaque white and one in transparent, in order to measure the anti-viral activity and cytotoxicity of a chemical compound in parallel ensuring the observed activity is not due to decreased cell proliferation.
- 10 *Renilla* luciferase HCV replicon cells (2209-23) cultured in Dulbecco's MEM (GibcoBRL cat no. 31966-021) with 5% fetal calf serum (FCS, GibcoBRL cat. no. 10106-169) were plated onto a 96-well plate at 5000 cells per well, and incubated overnight. Twenty-four hours later, different dilutions of chemical compounds in the growth medium were added to the cells, which were then further incubated at 37°C for three days. At the end of the
- 15 incubation time, the cells in white plates were harvested and luciferase activity was measured by using Dual-Luciferase reporter assay system (Promega cat no. E1960) All the reagents described in the following paragraph were included in the manufacturers kit, and the manufacturer's instructions were followed for preparations of the reagents. The cells were washed twice with 200 µl of phosphate buffered saline (pH 7.0) (PBS) per well and
- 20 lysed with 25 µl of 1x passive lysis buffer prior to incubation at room temperature for 20 min. One hundred microlitre of LAR II reagent was added to each well. The plate was then inserted into the LB 96V microplate luminometer (MicroLumatPlus, Berthold), and 100 µl of Stop & Glo® reagent was injected into each well and the signal measured using a 2-second delay, 10-second measurement program. IC₅₀, the concentration of the drug
- 25 required for reducing replicon level by 50% in relation to the untreated cell control value, can be calculated from the plot of percentage reduction of the luciferase activity vs. drug concentration.
- WST-1 reagent from Roche Diagnostic (cat no. 1644807) was used for the cytotoxicity assay. Ten microlitre of WST-1 reagent was added to each well including wells that
- 30 contain media alone as blanks. Cells were then incubated for 1 to 1.5 hours at 37 °C, and the OD value was measured by a 96-well plate reader at 450nm (reference filter at 650nm). Again CC₅₀, the concentration of the drug required for reducing cell proliferation by 50% in relation to the untreated cell control value, can be calculated from the plot of percentage reduction of the WST-1 value vs. drug concentration.

- 44 -

Compound Number	Luciferase Activity IC ₅₀ (μM)
5	5.33
18	2.4
25	2.47

EXAMPLE 9

Pharmaceutical compositions of the subject Compounds for administration via several routes were prepared as described in this Example.

5

Composition for Oral Administration (A)

Ingredient	% wt./wt.
Active ingredient	20.0%
Lactose	79.5%
Magnesium stearate	0.5%

The ingredients are mixed and dispensed into capsules containing about 100 mg each; one capsule would approximate a total daily dosage.

10

Composition for Oral Administration (B)

Ingredient	% wt./wt.
Active ingredient	20.0%
Magnesium stearate	0.5%
Crosscarmellose sodium	2.0%
Lactose	76.5%
PVP (polyvinylpyrrolidone)	1.0%

The ingredients are combined and granulated using a solvent such as methanol. The formulation is then dried and formed into tablets (containing about 20 mg of active compound) with an appropriate tablet machine.

15

Composition for Oral Administration (C)

Ingredient	% wt./wt.
Active compound	1.0 g
Fumaric acid	0.5 g
Sodium chloride	2.0 g
Methyl paraben	0.15 g
Propyl paraben	0.05 g
Granulated sugar	25.5 g
Sorbitol (70% solution)	12.85 g

- 45 -

Veegum K (Vanderbilt Co.)	1.0 g
Flavoring	0.035 ml
Colorings	0.5 mg
Distilled water	q.s. to 100 ml

The ingredients are mixed to form a suspension for oral administration.

Parenteral Formulation (D)

Ingredient	% wt./wt.
Active ingredient	0.25 g
Sodium Chloride	qs to make isotonic
Water for injection to	100 ml

5

The active ingredient is dissolved in a portion of the water for injection. A sufficient quantity of sodium chloride is then added with stirring to make the solution isotonic. The solution is made up to weight with the remainder of the water for injection, filtered through a 0.2 micron membrane filter and packaged under sterile conditions.

10

Suppository Formulation (E)

Ingredient	% wt./wt.
Active ingredient	1.0%
Polyethylene glycol 1000	74.5%
Polyethylene glycol 4000	24.5%

The ingredients are melted together and mixed on a steam bath, and poured into molds containing 2.5 g total weight.

15

Topical Formulation (F)

Ingredients	grams
Active compound	0.2-2
Span 60	2
Tween 60	2
Mineral oil	5
Petrolatum	10
Methyl paraben	0.15
Propyl paraben	0.05
BHA (butylated hydroxy anisole)	0.01
Water	q.s. 100

- 46 -

All of the ingredients, except water, are combined and heated to about 60°C with stirring. A sufficient quantity of water at about 60°C is then added with vigorous stirring to emulsify the ingredients, and water then added q.s. about 100 g.

5

Nasal Spray Formulations (G)

Several aqueous suspensions containing from about 0.025-0.5 percent active compound are prepared as nasal spray formulations. The formulations optionally contain inactive ingredients such as, for example, microcrystalline cellulose, sodium carboxymethylcellulose, dextrose, and the like. Hydrochloric acid may be added to adjust pH. The nasal spray formulations may be delivered via a nasal spray metered pump typically delivering about 50-100 microliters of formulation per actuation. A typical dosing schedule is 2-4 sprays every 4-12 hours.


The features disclosed in the foregoing description, or the following claims, or the accompanying drawings, expressed in their specific forms or in terms of a means for performing the disclosed function, or a method or process for attaining the disclosed result, as appropriate, may, separately, or in any combination of such features, be utilized for realizing the invention in diverse forms thereof.

The foregoing invention has been described in some detail by way of illustration and example, with reference to the specific embodiments for purposes of clarity and understanding. It will be obvious to one of skill in the art that changes and modifications may be made and equivalents substituted without departing from the true spirit and scope of the invention. Therefore, it is to be understood that the above description is intended to be illustrative and not restrictive. Many modifications may be made to adapt a particular situation, material, composition of matter, process, or process step or steps, to the objective spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto

All patents, patent applications and publications cited in this application are hereby incorporated by reference in their entirety for all purposes to the same extent as if each individual patent, patent application or publication were so individually denoted.

Claims

1. A compound of formula I



5 wherein:

R^1 and R^2 are independently selected from the group consisting of hydrogen, COR^5 , $C(=O)OR^5$, $C(=O)SR^5$, $C(=O)NHR^5$ and $COCH(R^6)NHR^7$;

R^3 and R^4 are independently selected from the group consisting of hydrogen COR^5 , $C(=O)OR^5$, $C(=O)SR^5$ and $COCH(R^6)NHR^7$, or R^3 and R^4 taken together are
10 selected from the group consisting of CH_2 , $C(CH_3)_2$ and $CHPh$;

R^5 is independently selected from the group consisting of C_{1-18} unbranched or branched alkyl, C_{1-18} unbranched or branched alkenyl, C_{1-18} unbranched or branched alkynyl, C_{1-18} lower haloalkyl, C_{3-8} cycloalkyl, alkyl substituted C_{3-8} cycloalkyl, phenyl optionally independently substituted with one to three
15 substituents selected from the group consisting of halo, lower alkyl, lower alkoxy, lower thioalkyl, lower alkyl sulfinyl, lower alkyl sulfonyl, nitro, and cyano, CH_2Ph wherein in phenyl ring is optionally substituted as described above and CH_2OPh wherein in phenyl ring is optionally substituted as described above;

20 R^6 is independently selected from the group consisting of the side chains of naturally occurring amino acids and C_{1-5} unbranched or branched alkyl; and

R^7 is selected from the group consisting of hydrogen, R^5OCO ; or,


R^6 and R^7 together are $(CH_2)_3$; and,

hydrates, solvates, clathrates and acid addition salts thereof; with the proviso that at least one of R^1 , R^2 , R^3 , or R^4 is other than hydrogen.

2. A compound according to claim 1 wherein R^1 , R^2 , R^3 , and R^4 each are independently COR^5 , $C(=O)OR^5$, $C(=O)SR^5$ and each R^5 is independently selected from the group consisting of C_{1-18} unbranched or branched lower alkyl, phenyl and CH_2OPh .
5
3. A compound according to claim 1 wherein R^1 , R^2 , R^3 , and R^4 are COR^5 and each R^5 is independently selected from the group consisting of C_{1-18} unbranched or branched lower alkyl, phenyl and CH_2OPh .
4. A compound according to claim 1 wherein R^1 is COR^5 , $C(=O)OR^5$, $C(=O)SR^5$ or
10 $COCH(R^6)NHR^7$ and R^2 , R^3 and R^4 are hydrogen.
5. A compound according to claim 4 wherein R^5 is selected from a group consisting of C_{1-18} unbranched or branched lower alkyl, C_{3-8} cycloalkyl, phenyl and CH_2OPh , or R^6 is selected from the group consisting of C_{1-5} unbranched or branched alkyl and the side chain of a naturally occurring amino acid.
- 15 6. A compound according to claim 1 wherein R^2 is selected from the group consisting of COR^5 , $C(=O)OR^5$, $C(=O)SR^5$, and $COCH(R^6)NHR^7$, R^1 , R^3 and R^4 are hydrogen.
7. A compound according to claim 6 wherein R^5 is selected from the group consisting of C_{1-18} unbranched or branched alkyl, C_{3-8} cycloalkyl and phenyl or R^6 is C_{1-5} unbranched or branched alkyl or the side chain of a naturally occurring amino acid.
- 20 8. A compound according to claim 6 wherein R^2 is $COCH(R^6)NH_2$ and R^6 is selected from the group consisting of C_{1-5} unbranched or branched alkyl and CH_2Ph .
9. A compound according to claim 1 wherein R^3 and R^4 both are hydrogen.
10. A compound according to claim 1 wherein R^1 is hydrogen and R^2 , R^3 and R^4 are independently selected from the group consisting of COR^5 , $C(=O)OR^5$ and
25 $C(=O)SR^5$.
11. A compound according to claim 1 wherein R^1 is hydrogen, R^2 is selected from the group consisting of COR^5 , $C(=O)OR^5$, $C(=O)SR^5$ and $COCH(R^6)NHR^7$, and R^3 and R^4 taken together are selected from the group consisting of CH_2 , $C(CH_3)_2$ and $CHPh$.

12. A compound according to claim 1 wherein R^1 and R^2 are hydrogen and R^3 and R^4 are independently selected from the group consisting of COR^5 , $C(=O)OR^5$, $C(=O)SR^5$ and $COCH(R^6)NHR^7$ wherein R^7 is hydrogen.
13. A compound according to claim 1 wherein R^1 and R^2 are independently selected from the group consisting of COR^5 , $C(=O)OR^5$, $C(=O)SR^5$ and $COCH(R^6)NHR^7$, and R^3 and R^4 taken together are selected from the group consisting of CH_2 , $C(CH_3)_2$ and $CHPh$.
14. A compound according to claim 1, wherein R^1 and R^2 are selected from the group consisting of hydrogen, COR^5 , $C(=O)OR^5$ and $COCH(R^6)NHR^7$; R^3 and R^4 independently of the other are selected from the group consisting of hydrogen, COR^5 , CO_2R^5 and $COCH(R^6)NHR^7$, or R^3 and R^4 taken together are selected from the group consisting of CH_2 , $C(CH_3)_2$ and $CHPh$; R^5 is independently selected from the group consisting of C_{1-6} unbranched or branched alkyl, C_{1-6} unbranched or branched alkenyl, C_{1-6} unbranched or branched alkynyl, C_{1-6} lower haloalkyl, C_{3-8} cycloalkyl, alkyl substituted C_{3-8} cycloalkyl, phenyl optionally independently substituted with one to three substituents selected from the group consisting of halo, lower alkyl, lower alkoxy, lower thioalkyl, lower alkyl sulfinyl, lower alkyl sulfonyl, nitro, and cyano, CH_2Ph wherein in phenyl ring is optionally substituted as described above and CH_2OPh wherein in phenyl ring is optionally substituted as described above; R^6 and R^7 are as defined in claim 1, with the proviso that at least one of R^1 , R^2 , R^3 , or R^4 is other than hydrogen.
15. A pharmaceutical composition comprising a therapeutically effective quantity of a compound as claimed in anyone of claims 1 to 14 and pharmaceutically acceptable excipients.
16. The use of a compound as claimed in anyone of claims 1 to 14 for the treatment of diseases.
17. The use of a compound as claimed in anyone of claims 1 to 14 for the manufacture of medicaments for the treatment of diseases mediated by the hepatitis C Virus (HCV).
18. A method for treating diseases mediated by the Hepatitis C Virus (HCV) virus comprising administering to a mammal in need thereof, a therapeutically effective quantity of a compound as claimed in anyone of claims 1 to 14.

- 50 -



wherein:

R¹ and R² are independently selected from the group consisting of hydrogen, COR⁵, C(=O)OR⁵, C(=O)SR⁵, C(=O)NHR⁵ and COCH(R⁶)NHR⁷;

5 R³ and R⁴ are independently selected from the group consisting of hydrogen, COR⁵, C(=O)OR⁵, C(=O)SR⁵ and COCH(R⁶)NHR⁷, or R³ and R⁴ taken together are selected from the group consisting of CH₂, C(CH₃)₂ and CHPh;

R⁵ is independently selected from the group consisting of C₁₋₁₈ unbranched or branched alkyl, C₁₋₁₈ unbranched or branched alkenyl, C₁₋₁₈ unbranched or branched alkynyl, C₁₋₁₈ lower haloalkyl, C₃₋₈ cycloalkyl, alkyl substituted C₃₋₈ cycloalkyl, phenyl optionally independently substituted with one to three substituents selected from the group consisting of halo, lower alkyl, lower alkoxy, lower thioalkyl, lower alkyl sulfinyl, lower alkyl sulfonyl, nitro, and cyano, CH₂Ph wherein in phenyl ring is optionally substituted as described above and CH₂OPh wherein in phenyl ring is optionally substituted as described above;

10

15

R⁶ is independently selected from the group consisting of the side chains of naturally occurring amino acids and C₁₋₅ unbranched or branched alkyl;


R⁷ is selected from the group consisting of hydrogen, R⁵OCO; or,

R⁶ and R⁷ together are (CH₂)₃; and,

20 hydrates, solvates, clathrates and acid addition salts thereof; with the proviso that at least one of R¹, R², R³, or R⁴ is other than hydrogen.

19. A process for converting an N-acyl cytidine compound IVa to a cytidine compound IVb by selective cleavage of an N-acyl moiety from IVa wherein:

- 51 -



R^2 , R^3 , R^4 and R^5 are as defined in claim 1 and

R^a is hydrogen or C_{1-4} alkyl;

said process comprising contacting a solution of said N-acyl pyrimidine nucleoside
 5 with ZnBr_2 in a protic solvent R^bOH wherein R^a is hydrogen or C_{1-4} alkyl.

20. A process according to claim 198 wherein said protic solvent is methanol.

21. A compound according to anyone of claims 1 to 14 whenever prepared according to claim 198.

22. The invention as described hereinbefore.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 03/12733

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07H19/167 A61K31/70 A61P31/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 02 100415 A (HOFFMANN LA ROCHE) 19 December 2002 (2002-12-19) page 16	1-22
A	WO 02 18404 A (HOFFMANN LA ROCHE) 7 March 2002 (2002-03-07)	
A	WO 93 17651 A (MAX DELBRUECK CENTRUM) 16 September 1993 (1993-09-16)	

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

26 March 2004

Date of mailing of the international search report

05/04/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bardili, W

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/12733

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 02100415	A	19-12-2002	CA 2449572 A1 WO 02100415 A2	19-12-2002 19-12-2002
WO 0218404	A	07-03-2002	AU 9549701 A BR 0113611 A CA 2419399 A1 CN 1466591 T WO 0218404 A2 EP 1315736 A2 US 2003008841 A1	13-03-2002 24-06-2003 07-03-2002 07-01-2004 07-03-2002 04-06-2003 09-01-2003
WO 9317651	A	16-09-1993	DE 4207363 A1 WO 9317651 A2	09-09-1993 16-09-1993